

APPROVED FOR RELEASE: 09/17/2001 , CIA-RDP86-00513R002201610009-0

AUTHOR:

TITLE:

PERIODICAL:

ABSTRACT:

SOV/19-59-7-158/369  
V.R., Mateyuk, N.A.,  
Jva, N.M.

Stabilization of a Single Twisted  
Natural or Artificial Fiber  
Byulleten' izobreteniy, 1959, Nr 7, pp 34-35 (USSR)  
Class 29a, 6.08. Nr 118945 (608873 of 3 October  
1958). The above device consists of a housing, To  
containing an upright pipe with hollow walls. To  
insure a uniform stabilization of a moving single  
thread in the steam-air medium and placed with  
machine, the pipe is funnel-shaped and filled with  
its large base, inside a container, supported by a plate,  
water. The container is supported by a plate,  
and connected with the angular source of feeding water  
by means of an angular pipe joint. A nickel-chro-  
mium heater is secured to the bottom of the con-  
tainer. The heater encloses a thread guide, fixed  
on the supporting plate. The upper end of the

SOV/19-59-4-162/317

14(1)

AUTHORS:

Kremnev, O.A., and Zozulya, N.V.

TITLE:

A Heating and Humidifying System

PERIODICAL:

Byulleten' izobreteniy, 1959, Nr 4, p 56 (USSR)

ABSTRACT:

Class 36d, 116. Nr 118340 (602192 of 20 June 1958).  
A system as in title, for heating and humidifying the air  
in a dwelling space, using a solution of lithium chloride  
as an intermediate heat carrier.

Card 1/1

"APPROVED FOR RELEASE: 09/17/2001

CIA-RDP86-00513R002201610009-0

APPROVED FOR RELEASE: 09/17/2001

CIA-RDP86-00513R002201610009-0"

KREMNEV, O. A.,

AP0049426

and red-hardness (for hardness HRC 60) 625-635°, i.e., values which are usually obtained for highly alloyed V-steels R014 and R14F4 with poorer grinding qualities. In steels with higher C, absence of ferrite, after heating to temps. above the pearlite transformation region, provides better dissoln. of carbides. In these steels pptn. of carbides during tempering takes place in a narrower temp. interval, hence the carbides are more dispersed and homogeneous in compn. and size, than in steels with lower C content. Increased C content to 1.0-1.05% is advisable for W-Mo steels, and partly also for R6M5 steel, particularly when they are used instead of highly alloyed V steels, because of their better mech. properties and grinding ability. Jiri Becvar

pc

42

19801263

1/2 007 UNCLASSIFIED PROCESSING DATE--13NOV70  
TITLE--INTENSIFICATION OF THE FINISHING OF A CUT STAPLE FIBER -U-

AUTHOR--(05)-KRENEV, O.A., BOROVSKIY, V.R., SATANDYSKIY, A.L., KIGEL,  
T.B., BUGAYCHOK, N.F.  
COUNTRY OF INFO--USSR

SOURCE--KHIM. VOLOKNA 1970, (2), 53-5 (RUSS)

DATE PUBLISHED-----70

SUBJECT AREAS--MATERIALS

TOPIC TAGS--RAYON, TEXTILE

CONTROL MARKING--NO RESTRICTIONS

DOCUMENT CLASS--UNCLASSIFIED

PROXY REEL/FRAE--2000/1661

STEP NO--UR/0183/70/000/002/0053/0055

CIRC ACCESSION NO--AP0125283

UNCLASSIFIED

2/2 007

UNCLASSIFIED

PROCESSING DATE--13NOV70

CIRC ACCESSION NO--AP0125283

ABSTRACT/EXTRACT--(U) GP-O- ABSTRACT. THE CON. CONTINUOUS WASHING OF RAYON STAPLE WITH DIL. NAOH SOLN. AT 78DEGREES REQUIRES 135 SEC TO REDUCE ITS S CONTENT BELOW 0.075PERCENT. THE INSERTION OF ROLLERS, WHICH COMPRESS THE STAPLE PERIODICALLY DURING ITS PASSAGE THROUGH THE BATH, AND TEMP. INCREASE TO 95DEGREES REDUCED THIS TIME TO 20-5 SEC. THE WASHING TIME WAS ALSO REDUCED TO 15-20 SEC WHEN VIBRATIONS OF 200 CYCLES-MIN AND 50 MM AMPLITUDE WERE APPLIED TO THE WASH SOLN. HELD AT 80DEGREES. BOTH OF THESE METHODS INCREASED PRODUCTIVITY. THE VIBRATIONS GAVE SOFTER, EASIER TO DRY FIBERS. FACILITY: INST. TEKHN. TEPLDIZ., KIEV, USSR.

UNCLASSIFIED

15(4)

SOV/19-59-7-354/369

AUTHORS: Kremnev, O.A., Borovskiy, V.R., Korostash, M.D.,  
Petrusha, G.I., Tyumenev, Ya.K., Burova, V.M.,  
Timofeyeva, Yu.A. and Zapol'skiy, I.A.

TITLE: A Method of Drying Cocoons

PERIODICAL: Byulleten' izobreteniy, 1959, No 7, pp 72-73 (USSR)

ABSTRACT: Class 82a, B. Nr 119141 (605185 of 1 August 1958).  
A method of drying cocoons, with the temperature  
of the drying agent falling in the drying process.  
The drying is carried out in a two-zone dryer and  
with artificial moistening of the drying agent be-  
ing fed into the chamber. In the first zone, the  
drying agent has temperature of 125°C and 25% humi-  
dity, and in the second zone, a temperature of 95°C  
and 35 to 40% humidity.

Card 1/1



15(4)

SOV/19-59-7-158/369

AUTHORS: Kremnev, O.A., Borovskiy, V.R., Mateyuk, N.A.,  
Filatov, G.M. and Antipova, N.M.

TITLE: A Device for Stabilization of a Single Twisted  
Thread, Made of Natural or Artificial Fiber

PERIODICAL: Byulleten' izobreteniy, 1959, Nr 7, pp 34-35 (USSR)

ABSTRACT: Class 29a, 6<sup>08</sup>. Nr 118945 (608873 of 3 October  
1958). The above device consists of a housing,  
containing an upright pipe with hollow walls. To  
insure a uniform stabilization of a moving single  
thread in the steam-air medium and on the twisting  
machine, the pipe is funnel-shaped and placed with  
its large base, inside a container, filled with  
water. The container is supported by a plate,  
and connected with the source of feeding water  
by means of an angular pipe joint. A nickel-chro-  
mium heater is secured to the bottom of the con-  
tainer. The heater encloses a thread guide, fixed  
on the supporting plate. The upper end of the

Card 1/2

SOV/19-59-7-158/369

A Device for Stabilization of a Single Twisted Thread, Made of  
Natural or Artificial Fiber

thread guide protrudes above the water level in  
order to prevent moistening of the thread. The  
lower end has its inlet enlarged for easing inser-  
tion of the thread.

Card 2/2

14(1)

SOW/19-59-4-162/517

AUTHORS: Krennev, C.A., and Zozulya, N.V.

TITLE: A Heating and Humidifying System

PERIODICAL: Byulleten' izobreteniy, 1959, Nr 4, p 36 (USSR)

ABSTRACT: Class 36d, 1<sub>16</sub>. Nr 118340 (602192 of 20 June 1958).  
A system as in title, for heating and humidifying the air  
in a dwelling space, using a solution of lithium chloride  
as an intermediate heat carrier.

Card 1/1

"APPROVED FOR RELEASE: 09/17/2001

CIA-RDP86-00513R002201610009-0



APPROVED FOR RELEASE: 09/17/2001

CIA-RDP86-00513R002201610009-0"

KREMNEV, O. A.,

1/2 028 UNCLASSIFIED PROCESSING DATE--20NOV70  
TITLE--DETERMINATION OF THE SOLUBILITY OF MOLYBDENUM CARBONYL IN CARBON  
MONOXIDE AT HIGH PRESSURE -U-  
AUTHOR--(04)--KRICHEVSKAYA, O.D., KREHNEV, V.L., ZELIKMAN, L.A., BOLOTOVA,  
K.N.  
COUNTRY OF INFO--USSR  
SOURCE--ZH. PRIKL. KHIM. (LENINGRAD) 1970, 43(4), 877-9  
DATE PUBLISHED--70  
SUBJECT AREAS--CHEMISTRY  
TOPIC TAGS--MOLYBDENUM COMPOUND, METAL CARBONYL, SOLUBILITY, CARBON  
MONOXIDE, HIGH PRESSURE  
CONTROL MARKING--NO RESTRICTIONS  
DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FRAME--3001/0439 STEP NO--UR/0080/70/043/004/0877/0879  
CIR ACCESSION NO--AP0126192

2/2 028 UNCLASSIFIED PROCESSING DATE--20NOV70  
CIRC ACCESSION NO--AP0126192  
ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. THE SOLY. WAS DETD. AT 100+280  
ATM. AND 0-30DEGREES. THE CONTENT OF MO(CO) SUB6 IN COMPRESSED CO ROSE  
WITH INCREASE IN PRESSURE. AN EXPRESSION IS GIVEN FOR THE SOLY. OF  
MO(CO) SUB6 IN CO AS A FUNCTION OF THE CO PRESSURE.

UNCLASSIFIED

USSR

UDC 547.341.26'118.07

LASKORIN, B. N., YAKSHIN, V. V., KREMNEVA, Ye. V., and SKOROVAROV, D. I.

"A Method of Making Monoethers of N-Substituted Thiocarbamoylphosphonic Acids"

Moscow, Otkrytiya, izobreteniya, promyshlennyye obraztsy, tovarnyye znaki, No 5, Feb 71, Author's Certificate No 292987, Division C, filed 11 Oct 69, published 15 Jan 71, p 101

Translation: This Author's Certificate introduces: 1. A method of making monoethers of N-substituted thiocarbamoylphosphonic acids. As a distinguishing feature of the patent, monosubstituted salts of monoalkylphosphites are interacted with alkyl- or arylisothiocyanates in the presence of heat in an organic solvent such as benzene, with subsequent treatment of the resultant product in a mineral acid such as hydrochloric acid, and isolation of the goal product by conventional methods. 2. A modification of this method distinguished by the fact that the process is carried out in the presence of a catalyst such as triethylamine.

1/1



USSR

UDC 547.341.26'118.07

LASKORIN, B. N., YAKSHIN, V. V., KREMNEVA, Ye. V., and SKORODVAREV, D. I.

"A Method of Making Monoethers of N-Substituted  $\alpha$ -Aminoalkylphosphonic Acids"

Moscow, Otkrytiya, izobreteniya, promyshlennyye obrabotsy, tovarnyye znaki, No 5, Feb 71, Author's Certificate No 292986, Class C, filed 11 Oct 69, published 15 Jan 71, p 101

Translation: This Author's Certificate introduces: 1. A method of making monoethers of N-substituted  $\alpha$ -aminoalkylphosphonic acids by interacting phosphites with Schiff bases in the presence of heat with subsequent treatment of the resultant product in a mineral acid such as hydrochloric acid. As a distinguishing feature of the patent, the process is simplified by using monosubstituted ammonium, alkali or alkaline earth salts of monoalkylphosphites as the phosphite, and carrying out the process in the presence of a catalyst such as triethylamine. 2. A modification of this method distinguished by the fact that heating is done to a temperature of 110-120°C.

1/1

- 18 -

Acc. Nr: **AP0052066**

K

Ref. Code: **UR0396**

PRIMARY SOURCE: Patologicheskaya Fiziologiya i  
Eksperimental'naya Terapiya, 1970, Vol 14,  
Nr 1, pp 29-33

OXYGEN TENSION IN THE WALL OF THE STOMACH, SMALL AND LARGE  
INTESTINE UNDER NORMAL AND PATHOLOGICAL CONDITIONS

A. P. Krandal', I. M. Epshteyn, S. V. Yeliseyeva

A total of 36 acute experiments were carried out on cats. Using a copper amalgamated electrode  $pO_2$  was determined in the wall of the stomach, small and large intestine. The highest initial  $pO_2$  level was found in the wall of the small intestine, and the lowest -- in the large intestine, the stomach occupying an intermediary position. Under pathological conditions (hyper- and hypoxia, hypervolemia, ischemia) alteration of  $pO_2$  level in the wall of the small intestine considerably exceeded these changes in the wall of the stomach and large intestine. The fall of  $pO_2$  caused by blood letting and ligation of the vessels can be compensated by giving pure oxygen to inhale. Different conditions of oxygen supply of the stomach and intestine are apparently conditioned by peculiarities attending the blood supply and the metabolism of these organs.

111

44 2

REEL / FRAME  
**19820604**

USSR

UDC: 681.3

BOYKOV, V. N., ~~SEENDEL~~ Yu. M., RABINOVICH, V. I., TROFIMOV, G. Ye.,  
SHUL'TS, V. P.

"On Calculating the Precision of Digital Measuring Instruments"

V sb. Teoriya i praktika ispol'z. sredstv tekhn. kibernetiki. Kn. 1 (Theory and Practice in Utilizing the Facilities of Technical Cybernetics--collection of works. Book 1), Novosibirsk, 1970(1971), pp 94-99 (from EZh-Matematika, No 11, Nov 71, Abstract No 11V786)

Translation: The authors consider the effect which improper operation of flip-flops and comparison devices has on the accuracy of a digital device based on the method of binary sweep balancing. In order to explain "in pure form" the nature of the effect which these elements have on the operating precision of the device, a separate analysis is made: for instance, in studying the influence of incorrect flip-flop operation, the comparison device is first assumed as ideal, then real. For the sake of definiteness, distribution of the measured quantity  $X$  is assumed to be uniform, and the average modulus of the error is taken as the quality criterion for accuracy of the device. It is noted that all results can be recorded in general form or obtained in the same way for specific distributions of  $X$  with the use of many other metrological criteria; e. g., variance or mathematical expectation of the error, etc. V. Mikhayev.  
1/1

- 30 -

USSR

UDC: 681.3

BOYKOV, V. N., KRENDEL', Yu. M., RABINOVICH, V. I., TROFIMOV, O. Ye.,  
SHUL'TS, V. P.

"On Calculating the Accuracy of Digital Measuring Instruments"

V sb. Teoriya i praktika ispol'z. sredstv tekhn. kibernetiki. Kn. 1 (Theory  
and Practice in Using the Facilities of Technical Cybernetics. Book 1--  
collection of works), Novosibirsk, 1970(1971), pp 94-99 (from RZh-Kibernetika,  
No 11, Nov 71, Abstract No 11V786)

Translation: The authors consider the effect which improper operation of  
the flip-flops and the comparison unit has on the accuracy of a digital  
device based on the method of binary sweep balancing. In order to explain  
the nature of the effect which these elements "in pure form" have on the  
accuracy of the device, a separate analysis is made: for instance in  
studying the effect of improper operation of the flip-flops, the comparison  
unit is first assumed to be ideal, and then real. It is assumed for the  
sake of definiteness that the distribution  $X$  of the measured quantity is  
uniform, and the average absolute value of the error is used as the cri-

1/2

USSR

BOYKOV, V. N. et al., Teoriya i praktika ispol'z. sredstv tekhn. kibernetiki.  
Kn. 1, Novosibirsk, 1970(1971), pp 94-99

terion of accuracy of the instrument. It is noted that all the results may be written in general form or derived in the same way for distributions X using many other metrological criteria (variance of the error, mathematical expectation of the error, etc.). V. Mikheyev.

2/2

- 17 -



KRENKE, A.N.

Glaciology

GLACIOLOGICAL ADVANCES OF GLACIOL

Article by Doctor of Geography Sciences L. N. Krenke, Candidate of Geographic Sciences A. N. Krenke, and Candidate of Geographic Sciences V. N. Krenke, Moscow, 1977

The rapid advance of glaciers, observed since the beginning of the 20th century, has become a natural phenomenon stimulated by instability of the glacial system. This is based on the possibility of predicting the future of the glacial system. The possibility of predicting the future of the glacial system is based on the possibility of predicting the future of the glacial system. The possibility of predicting the future of the glacial system is based on the possibility of predicting the future of the glacial system.

Investigations in recent decades, performed both in the USSR and abroad, have established that periodic advances of glaciers are a natural phenomenon stimulated by instability of the glacial system. This is based on the possibility of predicting the future of the glacial system. The possibility of predicting the future of the glacial system is based on the possibility of predicting the future of the glacial system. The possibility of predicting the future of the glacial system is based on the possibility of predicting the future of the glacial system.

A study of glaciological data of the Institute of Geography of the USSR Academy of Sciences, Moscow, has shown that the advance of glaciers is a natural phenomenon stimulated by instability of the glacial system. This is based on the possibility of predicting the future of the glacial system. The possibility of predicting the future of the glacial system is based on the possibility of predicting the future of the glacial system.





USSR UDC 543.422.4:542.957:547.357.4:546.821'131:547.558.1:547.315.1

BONDARENKO, G. N., KHAR'KOVA, Ye. M., MUSHINA, Ye. A., ~~TERENINA, M. P.~~, and  
KRENTSEL', B. A., Institute of Petrochemical Synthesis Imeni A. V. Topchiyev,  
Academy of Sciences USSR

"IR Spectra of the Reaction Products of Bis- $\pi$ -allylnickel Bromide With  
Titanium Tetrachloride, Triphenylphosphine, and Allene"

Moscow, Izvestiya Akademii Nauk SSSR, Seriya Khimicheskaya, No 6, Jun 73,  
pp 1224-1227

Abstract: The IR spectra were studied of the reaction products of bis- $\pi$ -  
allylnickel bromide with  $\text{TiCl}_4$  and triphenylphosphine. It was proposed  
that there occurs a  $\pi \rightarrow \sigma$  isomerization in  $\pi$ -allylnickel complex. The  
 $\pi$ -allyl structure is retained in the reaction of bis- $\pi$ -allylnickel bromide  
with allene. During allene polymerization the activity of the complexes  
decreases in the order  $(\pi\text{-C}_3\text{H}_5\text{Ni})_2 > (\pi\text{-C}_3\text{H}_5\text{NiBr})_2 > (\pi\text{-C}_4\text{H}_7\text{Cl})_2$ .

1/1

KRENTSEV

BA

Chemistry

SYNTHESIS OF ACTIVE SYNTHETIC POLYMERS AND MACROMOLECULAR MODELS OF

Article by Doctor of Chemical Sciences B. A. Krentsev, Moscow, Vostochny  
Akademik Nauk SSSR, Russian, Vol. 62, No. 4, April 1977, pp. 117-119

Synopsis in English

The development of polymer science has entered a revolutionary stage in-  
fluence on the natural sciences as a whole, especially on their physico-  
chemical, biochemical, and biological aspects. New concepts about the crea-  
tion of physiologically active polymeric substances differing from their  
low-molecular analogs in their action on living organisms were a reflection  
of that influence.

Symposia on the creation of physiologically active polymeric substan-  
ces and study of their properties have become traditional. Two such sym-  
posia were held in Riga in 1962 and 1963. On 16-19 November 1971 a third sym-  
posium was held at the same place, and when differed from preceding ones  
by the fact that its program included a new and promising direction -- the  
development and investigation of macromolecular models of signaling. The  
expansion of the third symposium also was reflected in the compo-  
sition of its participants, among them were organic chemists, physical chem-  
ists, biologists, and specialists in medicine, biochemists -- a total of about 200 per-  
sons from various scientific institutions of Moscow, Leningrad, Riga, Kiev  
and other cities of the country.


The primary reports were heard. V. A. Krentsev arrived in Riga at  
the first symposiums in the use of polymeric substances as carriers of  
biological activity. He held expert-panel discussions showing that the creation  
of a medicinal substance with a polymer carrier permits not only increas-  
ing the effect of its action but also discovering new medical functions.  
The results of the report aroused objections -- for example, the advan-  
tages of creating "polymeric" antiparasitics. Evidently in that case there  
will be no sufficiently reliable experimental data for unequivocal decisions  
and further experimental work is required. The reporter and the participants  
of the symposium who spoke in the discussion turned attention toward the main

Acc. Nr. **AP0100204**

Abstracting Service:  
CHEMICAL ABST. 6-70

Ref. Code  
**UR0062**

/ 111851) Decarboxylation of polymeric acids with conjugated bonds. Yandarova, M. N.; Geiderikh, M. A.; Bepitsky, B. A. (Inst. Neftekhim. Sin. im. Topchieva, Moscow, USSR). *Izv. Akad. Nauk SSSR, Ser. Khim.* 1970, (1), 75-84 (Rus). The irradiation of  $\text{HC}\equiv\text{CCO}_2\text{H}$  (I) in the solid phase gives *trans,trans*-poly(propionic acid) (II); in the liq. phase, *trans,cis* (III) and *cis,trans*-poly(propionic acid) (IIIa) are obtained (G. V. Khutareva, et al., 1965). The decarboxylation of II, III, IIIa.



poly(acrylic acid) (IV) and 1-H<sub>2</sub>C:CHCO<sub>2</sub>H copolymers was compared. The conjugation reduces the decarboxylation activation energy of II, III, and IIIa to 6-12.5 kcal/mole as compared with 42.7 kcal/mole for IV. The configuration also affects the decarboxylation rates; III in which there is no H bonding between CO<sub>2</sub>H groups is decarboxylated faster than II or IIIa. (P.H.)

REEL/FRAME  
**19841586**

106

Acc. Nr:

AP0049793

Abstracting Service:

CHEMICAL ABST. 5-70

Ref. Code:

UR 0183

101698a Copolymerization of vinyl chloride with acrylic acid. Ruchinski, S. P.; Zay'nylov, A. N.; Krut'nyy, B. K.; Krentsel, B. A. (USSR). *Khim. Volokna* 1970, (1), 75 (Russ). Vinyl chloride (I) and acrylic acid (II) were copolymerized in MeOH contg. some H<sub>2</sub>O at atm. pressure and -10 to 5° with an initiating system comprising (NH<sub>4</sub>)<sub>2</sub>S<sub>2</sub>O<sub>8</sub>, Rongalite, and Cu salts to give I-II copolymer (III), which was more coned. in II than the original monomer mixt., dyed well with basic dyes, had glass transition temp. ~100°, was more sol. in org. solvents than pure I polymer (IV) and formed fibers from acetone and HCONMe<sub>2</sub> solns. Fibers of III having 10% and 27.2% II had lower tex. strength, and % elongation than IV fibers, but higher shrink resistance (except for III with 27.2% II heated in air at 130°).

BCJR 1

28.

REEL/FRAME  
19801715

USSR

UDC 669.01:536.425:536.2.001.2

ZINOV'YEV, V. YE., ~~KRECHTSIS, P. P.~~, GEL'D, P. V.

"Effect of Phase Transformation and Thermal Defects on the Thermal Conductivity and Thermal Diffusivity of Ti, Fe, Co, Ni, Pt and certain Steels at High Temperatures"

V sb. Fiz.-khim. osnovy proiz-va stali (Physical-Chemical Principles of Steel Production -- collection of works), Moscow, Nauka Press, 1971, pp 91-96 (from RZh--Metrologiya i Izmeritel'naya Tekhnika, No 3, Mar 72, Abstract No 3.32.979)

Translation: The block diagram is presented for a device on which the method of plane temperature waves is used to measure the coefficient of thermal diffusivity in thin plane-parallel plates of metals and alloys at temperatures above 700° C. Results are presented for measuring the coefficients of thermal diffusivity of Ti, Fe, Co, Ni, Pt and EI929, EI826, EI47B3 and EI435 steel. The thermal conductivity of the given metals and alloys was calculated using the published data on the density and heat capacity. There are 4 illustrations and a 27-entry bibliography.

1/1

USSR

UDC: 669.018.5:536.41

KRENTSIS, R. P., KALISHEVICH, G. I., GEL'D, P. V., and ANDRIENKO, L. P.

"Thermal Expansion of Chromium, Manganese, Iron, and Cobalt Silicides"

Tomsk, Izvestiya VUZ--Fizika, No 1, 1972, pp 153-155

Abstract: This brief communication gives the results of measurements made by the authors of the coefficients of linear expansion of various specimens as a function of heat in a temperature range of 100 to 1000° K. The specimens were chromium, manganese, iron, and cobalt silicides having the same cubic structure of the B20 type. The specimens were made of 99.98% Cr by weight; 99.95% Mn; 99.95% Fe; 99.98 Co; and monocrystalline silicon, 99.997% Si in an induction furnace of the MVP-3M type, in an argon atmosphere. After alloying, the specimens were sucked up in quartz capillary tubes 3-4 mm in diameter. The errors made in the measurements did not exceed 2-3%, except for the lowest temperature region of 100-200° K, when the error amounted to 4-5%. Curves are given for the expansion coefficients as functions of the temperature, and a table giving some characteristics of the four types of monosilicides is presented. The authors are connected with the S. I. Kirov Ural Polytechnical Institute.

1/1

1/2 023 UNCLASSIFIED PROCESSING DATE--18SEP70  
TITLE--THERMAL CONDUCTIVITY AND THERMAL DIFFUSIVITY OF IRON MONOSILICIDE  
-U-  
AUTHOR--(03)-KRENTSIS, R.P., OSTROVSKIY, F.I., GELD, P.V.  
COUNTRY OF INFO--USSR *R*  
SOURCE--FIZ. TEKH. POLUPROV. 1970, 4(2), 403-5  
DATE PUBLISHED-----70  
  
SUBJECT AREAS--PHYSICS, CHEMISTRY  
TOPIC TAGS--IRON COMPOUND, THERMAL CONDUCTIVITY, SILICON COMPOUND,  
ELECTRICAL CONDUCTIVITY, SILICON SEMICONDUCTOR  
  
CONTROL MARKING--NO RESTRICTIONS  
DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FRAME--1988/0573 STEP NO--UR/0449/00/004/002/0403/0405  
CIRC ACCESSION NO--AP0105558  
UNCLASSIFIED

2/2 023 UNCLASSIFIED PROCESSING DATE--18SEP70  
 CIRC ACCESSION NO--AP0105558  
 ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. SINGLE CRYSTALS OF FESI WERE GROWN FROM THE MELT BY THE CZOCHRALSKI METHOD, THE FESI BEING PREPD. FROM CARBONYL FE AND SEMICONDUCTOR GRADE SI. THE THERMAL COND. WAS MEASURED BY USING EQUIPMENT DESCRIBED PREVIOUSLY (L. P. FILIPPOV, 1967) OVER THE TEMP. RANGE 400-900DEGREES, WHILE THE ELEC. RESISTANCE WAS MEASURED POTENTIOMETRICALLY. AN EQUATION IS GIVEN RELATING THE ELEC. RESISTANCE AND THE THERMAL COND. THIS IS THEN USED TO CALCULATE THE LORENZ NO. FOR FESI OVER THE TEMP. RANGE 1200-1600DEGREES WHERE THE ELEC. RESISTANCE IS ALMOST CONST. THE VALUES OBTAINED WERE  $L_{SUBMIN}$ , EQUALS  $3.1 \times 10^{-8}$  PRIME NEGATIVE 8 V PRIME 2-DEGREE PRIME 2 AND  $L_{SUBMAX}$ , EQUALS  $3.4 \times 10^{-8}$  PRIME NEGATIVE 8 V SUB 2, DEGREE PRIME 2, WHICH ARE SIGNIFICANTLY HIGHER THAN THE VALUE FOR METALS, I. E.  $2.45 \times 10^{-8}$  PRIME NEGATIVE 8 V PRIME 2-DEGREE PRIME 2.

UNCLASSIFIED



1/2 026 UNCLASSIFIED PROCESSING DATE--18SEP70  
TITLE--ANISTROPY OF THE THERMAL EXPANSION OF IRON DIGERMANIDE -U-

AUTHOR--(03)-KRENTSIS, R.P., MIKHELSON, A.V., GELD, P.V.

COUNTRY OF INFO--USSR

SOURCE--FIZ. TVERD. TELA 1970, 12(3), 933-4

DATE PUBLISHED-----70

SUBJECT AREAS--MATERIALS, PHYSICS

TOPIC TAGS--PHYSICAL CHEMISTRY PROPERTY, SINGLE CRYSTAL, IRON COMPOUND,  
GERMANIUM COMPOUND, IRON COMPOUND, THERMAL EXPANSION, PHASE TRANSITION

CONTROL MARKING--NO RESTRICTIONS

DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FRAE--1988/0583

STEP NO--UR/0181/70/012/003/0933/0934

CIRC ACCESSION NO--AP0105566

UNCLASSIFIED

2/2 026

UNCLASSIFIED

PROCESSING DATE--19 SEP 70

CIRC ACCESSION NO--AP0105566

ABSTRACT/EXTRACT--(U) GP-O- ABSTRACT. THE LINEAR EXPANSION COEFF. WAS MEASURED AT 90-480 DEGREES K BY USING 2 SPECIMENS CUT FROM A SINGLE CRYSTAL OF FE<sub>2</sub> SUB<sub>2</sub> IN THE DIRECTIONS (001) AND (100). THE DATA OBTAINED ARE GENERALIZED GRAPHICALLY AND THEY INDICATE A CONSIDERABLE ANISOTROPY OF THE PHYS. PROPERTIES OF THE COMPD. FE<sub>2</sub> SUB<sub>2</sub> AND SHARP ANOMALIES OF THEIR TEMP. DEPENDENCES IN THE REGION OF THE TRANSFORMATION. DIFFERENCE IN SIGN AND THE MAGNITUDE OF THE ANOMALIES ON THE CURVES ALPHA (TAU) ARE APPARENTLY RELATED TO THE DIFFERENCES IN THE CHARACTER OF THE EXCHANGE INTERACTION BETWEEN THE ATOMS LOCATED IN THE SAME AND NEIGHBORING BASIS PLANES OF THE TETRAGONAL STRUCTURE OF FE<sub>2</sub> SUB<sub>2</sub>. THE TEMP. SHIFT OF THE NEEL TEMP. CAN BE ATTRIBUTED EITHER TO BROADENING OF THE TRANSFORMATION OR TO THE PRESENCE OF A SERIES OF SUCCESSIVE PHASE TRANSITIONS.

UNCLASSIFIED

USSR

K UDC (546.72'28 + 546.73'28):548.55

FROLOV, A. A., ~~KRENTSIS, R. P.~~, and GEL'D, P. V., Ural Polytechnic Institute imeni S. M. Kirov, Sverdlovsk, Ministry of Higher and Secondary Specialized Education RSFSR  
"Growth of FeSi and CoSi Single Crystals by Czochralski Method"

Moscow, Izvestiya Akademii Nauk SSSR -- Neorganicheskiye Materialy, Vol 6, No 4, Apr 70, pp 828-829

Abstract: The article describes the growth of FeSi and CoSi single crystals by the Czochralski method. The starting materials were K-O brand cobalt, V-3 brand carbonyl iron and single-crystal silicon. Single crystals were grown from the melt on a ZhK 01.01 device (with graphite heater) in a vacuum of  $5 \cdot 10^{-5}$  mm Hg. Pulling rate 0.5-0.75 mm/min. The crucible and seed rotated in opposite directions at 30 and 45 rpm respectively. In some cases the upper rod remained immobile. The crucibles were of  $Al_2O_3$ . The composition of the grown single crystals was determined by X-ray from lattice parameters. The microhard-

1/2

USSR

FROLOV, A. A., et al., Izvestiya Akademii Nauk SSSR -- Neorganicheskiye Materialy, Vol 6, No 4, Apr 70, pp 828-829

ness of unetched sections was measured on a PMT-3 device and was found to remain constant according to ingot length, which indicates the mechanical homogeneity of the single crystal.

2/2

1/2 033 UNCLASSIFIED PROCESSING DATE--30OCT70  
TITLE--THERMAL DIFFUSIVITY AND THERMAL CONDUCTIVITY OF SOLID SOLUTIONS OF  
SILICON IN IRON AND COBALT -U-  
AUTHOR-(04)-KRENTSIS, R.P., ZINOVYEV, V.YE., ANDREYEVA, L.P., GELD, P.V.  
COUNTRY OF INFO--USSR  
SOURCE--FIZ. METAL. METALLOVED. 1970, 29(1), 118-23  
DATE PUBLISHED-----70  
SUBJECT AREAS--MATERIALS, PHYSICS  
TOPIC TAGS--THERMAL CONDUCTIVITY, SOLID SOLUTION, IRON ALLOY, COBALT  
ALLOY, SILICON CONTAINING ALLOY, CURIE POINT, THERMAL DIFFUSION, HIGH  
PURITY METAL, PHONON SCATTERING  
CONTROL MARKING--NO RESTRICTIONS  
DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FAME--1998/0941 STEP NO--UR/0126/70/029/001/0118/0123  
CIRC ACCESSION NO--AP0121543  
UNCLASSIFIED

2/2 033

UNCLASSIFIED

PROCESSING DATE--300CT70

CIRC ACCESSION NO--AP0121543

ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. THE THERMAL DIFFUSIVITY OF A SOLID SOLN. OF SI IN FE (UP TO 8.2 WT. PERCENT SI), OF SI IN CO (UP TO 3 WT. PERCENT SI), AND OF PURE CO WAS DETD. IN THE REGION OF THE CURIE TEMP. (T SUBC). IN EACH CASE, THE POLYTHERM OF THERMAL DIFFUSIVITY SHOWS A MIN. IN THE REGION OF T SUBC. AWAY FROM T SUBC, AN APPROX. LOGARITHMIC DEPENDENCE WAS OBSD. OF THERMAL DIFFUSIVITY ON T-T SUBC. THE DEPENDENCE IS MORE COMPLEX CLOSE TO T SUBC. A MIN. IN THE THERMAL COND. POLYTHERMS AT T SUBC ARE A RESULT OF A STRONG PHONON ELECTRON AND PHONON SPIN SCATTERING. THE MECHANISM OF SCATTERING AND TRANSFER IN FERRONAGNETIC SYSTEM IS DISCUSSED. FACILITY: URAL. POLITEKH. INST.IM. KIROVA, SVERDLOVSK, USSR.

UNCLASSIFIED

USSR

UDC 621.371.332.4

SEMAKOV, V. L., KREPAK, V. N., and YAKIMENKO, I. P.

"Scattering of Electromagnetic Waves by Cylindrical Systems With Heterogeneous Gyrotropic Plasmas"

Moscow, V sb. X Vses. konf. po rasprostr. radiovoln. Tezisy dokl. Sekts. 5 (Tenth All-Union Conference on the Propagation of Radio Waves; Report Theses; Section 5--collection of works) "Nauka," 1972, pp 22-26 (from RZh--Radiotekhnika, No 10, 1972, Abstract No 10A377)

Translation: A solution is found for the problem of the scattering of a plane wave by cylindrical objects with radially heterogeneous gyrotropic plasmas. The effect of an external magnetic field, the frequency, and the nature of the radial variation in the electron density of the plasma, is investigated. Six illustrations, bibliography of four. N. S.

1/1

KREP'KAYA, V.P.

59213  
6-73

2-1. DISTRIBUTION OF ADVERTISEMENTS IN THE EPITAXIAL LAYERS OF CALCIUM PHOSPHIDE  
MIXED WITH SULFUR AND NITROGEN

Article by G. M. Izrael, A. N. Petrov, A. N. Plent, D. A. Yakovlev, V. P. Krep'kaya, V. V. Artyukova, Leningrad: Novosibirsk, Ill. Sibirskiy na Prots-  
essakh Natsionalnaya Poluprovodnikovaya Elektronika, Moscow, 1973, 1-17  
June 1974, 5-111

The properties of monocristalline layers of calcium phosphide alloyed  
with nitrogen were investigated. The layers were grown by the method of liquid  
phase epitaxy in an open system on GaP substrates alloyed with sulfur obtained  
by the electrochemical method. Alloying with nitrogen was realized by adding sel-  
enium nitride from 0.05 to 1 percent by weight to the GaP-Ge melt. The sulfur  
goes into the melt as a result of solution of the substrate.

By using the measurements of the capacitance of the surface barrier at  
the GaP-As contact, a study was made of the concentration distribution of the  
donors with respect to depth of the epitaxial layer. On the film obtained,  
a study was made of the optical absorption near the edge of the basic band.  
On the basis of an analysis of the absorption spectra, the nitrogen concentra-  
tion in the film was determined as a function of the amount of GaP in the Ga-  
Ge melt. Results are presented from a study of the electroluminescence of the  
p-n junctions obtained by the diffusion of zinc in the arsenic layers. The radi-  
ation spectra at room temperature contained two bands with 2.16 electron volt  
and 1.12 electron volt peaks.

The quantum yield of green radiation is  $4 \cdot 10^{-3}$ . The brightness of the  
diodes at a current density of 10 milliamperes/cm<sup>2</sup> was 30 mcd at 100 milliwatts/  
cm<sup>2</sup>; it was 1,100 mcd.



USSR

BELOUSOVA, I. M., BOBROV, B. D., KITSELEV, V. M., KURZENKOV, V. N., KREPOSTNOV, P. I.

"Photodissociative  $I^{127}$  Laser in a Magnetic Field"

Moscow, Zhurnal Eksperimental'noy i Teoreticheskoy fiziki, Vol 65, No 2 (8), 1973, pp 524-536

Abstract: A study was made of the effect of a magnetic field and also a number of other factors on the kinetics of the radiation spectrum of a photodissociative  $I^{127}$  laser in the  $2P_{1/2} - 2P_{3/2}$  transition. The behavior of the spectrum of the induced radiation was investigated in the presence and absence of a magnetic field, and the superthin splitting constant of the upper operating level of the iodine atom  $A_{1/2}$  was determined experimentally. A broadening of the luminescence line in the operating transition of the iodine atom during collisions with  $C_2F_4$  molecules and also with argon and xenon atoms found. Then the corresponding broadening cross sections and the Van der Waals constants were determined for the interaction of the iodine atom with these gases. The calculated values of the frequencies and amplification factors for the most intense groups of Zeeman components were obtained with a varia-

USSR

BELOUSOVA, I. M., et al., Zhurnal Eksperimental'noy i Teoreticheskoy Fiziki, Vol 65, No 2 (8), 1973, pp 524-536

tion explaining the behavior of the induced radiation spectrum with the variation of the magnetic field. The magnitude of the relaxation between sublevels with different  $F$  of the upper operating state  $^2F_{1/2}$  was evaluated.

The study of the spectral composition of the induced radiation in the  $^2P_{1/2} - ^2P_{3/2}$  transition of the iodine atom showed that the radiation kinetics of the given laser, which is a complex spectral system, depends on a number of factors such as the magnitude of the magnetic field, the gas pressure in the cell, and the magnitude of the pumping energy. The characteristics of the output radiation of the laser are determined not only by the kinetics of the chemical reactions but also by the structure of the upper and lower transition states, which varies even during the oscillation pulse time. This factor must be considered when constructing various kinetic models of the photodissociative iodine laser. In turn, for a more detailed study of the behavior of the oscillation spectrum in a magnetic field the population kinetics on all sublevels of the investigated transition must be considered.

2/2

USSR

UDC: 620.279.16

NIKIFORENKO, Zh. G., SYSKOV, V. A., KREPS, N. S., IGHNATINSKIY, I. L.,  
VNIINK, Kishinev

"Ultrasonic Inspection of Bilayer Plates"

Sverdlovsk, Defektoskopiya, No 3, May/Jun 71, pp 87-93

Abstract: In order to evaluate the possibility of inspecting bilayer plates by the ultrasonic reflected pulse resonance method (L. G. Merkulov, V. M. Verevkin, Defektoskopiya, 1965, No 5, p 13) when the contacting liquid can wet only one surface of the plate to feed ultrasonic oscillations into the material being inspected, the authors calculate the shape of a pulse reflected from a plate separating liquid from air, disregarding attenuation. A block diagram of the ultrasonic flaw detector is presented. The proposed device can be used to check for diffusion welds between the layers of double-layer plates. The results of experiments show that the Merkulov-Verevkin method is a sound basis for highly sensitive high-productivity flaw detection.

1/1

USSR

KREPS, V. L.

"Finite Non-coalition Games with Dependent Strategies"

Teoriya Igr [Games Theory -- Collection of Works], Yerevan, 1973,  
pp 211-215 (Translated from Referativnyy Zhurnal Kibernetika, No 10,  
1973, Abstract No 10V457)

Translation: The type of dependence of the strategies of all players  
is characterized by real function  $\phi$ , which gives the rule for production  
of a measure in a set of situations from measures in sets of strategies.  
In case of independence, this function is a product. The relationship  
between the existence of the situation of equilibrium and the type of  
dependence is studied.

L. Brublevskaya

1/1

USSR

UDC 612.822.1+612.82

KREPS, Ye. M., Academician, Institute of Evolutionary Physiology and Biochemistry imeni I. M. Sechenov, USSR Academy of Sciences, Leningrad

"Functional Biochemical Changes in the Rat Brain During the Initial Stage of the Effects of Increased Oxygen Pressure"

Moscow, Doklady Akademii Nauk SSSR, Vol 197, No 2, 1971, pp 494-496

Abstract: The effects of oxygen inhaled under high pressure are usually divided into three stages: the initial, or compensatory stage, the preconvulsive, or forerunner stage, and the convulsive or agonal stage. In this investigation, the behavior of rats, the cholinesterase activity in the cerebral hemispheres, and the concentration of ascorbic acid in the adrenal glands were studied in the initial stage of oxygen toxicity. The tests were done on rats before and after they were kept in a hyperbaric chamber filled with pure oxygen at 2.5 atm. After the 30-40 minute exposure, the orientation and search activity of the rats increased 63%, cholinesterase activity increased 10%, and ascorbic acid concentration decreased 24%. The conclusion was drawn that the initial stage of oxygen toxicity involves not only compensatory mechanisms but also a general sympathetic excitation representing a nonspecific reaction to stress.

1/1

USSR

KREPS, Ye. M., Academician (Editor)

Mekhanizmy regulyatsii fiziologicheskikh funktsiy, Sbornik statey k 40-letiyu nauchnoy deyatel'nosti akademika Vladimira Nikolayevicha Chernigovskogo (Regulatory Mechanisms of Physiological Functions -- Collection of Articles for the 40th Anniversary of the Scientific Activity of Academician Vladimir Nikolayevich Chernigovskiy), Nauka Press, Leningrad, 1971, 316 pp

Translation: Contents:  
Foreword

Page  
3

General Problems of Physiology

|                                                                                                                               |    |
|-------------------------------------------------------------------------------------------------------------------------------|----|
| E. A. Asratyan, Role of Inhibition in the Fatigue Phenomenon                                                                  | 6  |
| D. A. Biryukov, (Deceased) and V. I. Klimova-Cherkasova, Central Tonic Mechanisms of the Regulatory Functions of the Brain    | 8  |
| P. G. Borach, Effect of Thermal, Osmotic and Chemical Stimulation of the Hypothalamus on Food and Water Intake                | 16 |
| I. Ye. Ganelina, M. Ya. Ratner, and A. Ya. Yaroshevskiy, (Deceased) Clinical Observations as Method of Research in Physiology | 24 |
| G. V. Gershuni, Some Features of Brain Activity                                                                               | 43 |
| V. S. Gurfinkel' and M. L. Shik, Multilevel Motor Control                                                                     | 48 |

1/5

USSR

KREPS, Ye. M., Mekhanizmy regulyatsii fiziologicheskikh funktsiy, Sbornik statey k 40-letiyu nauchnoy deyatel'nosti akademika Vladimira Nikolayevicha Chernigovskogo, Nauka Press, Leningrad, 1971, 316 pp

|                                                                                                                                |            |
|--------------------------------------------------------------------------------------------------------------------------------|------------|
| V. V. Zakusov and N. V. Kaverina, Pharmacological Effects on the Central Adrenergic Mechanisms of Regulating Blood Circulation | Page<br>55 |
| G. P. Konradi and L. A. Vil'de, Reactions of the Capacitive Vessels                                                            | 63         |
| P. G. Kostyuk and A. I. Pilyavskiy, Neuron Organization of the Rubrospinal System                                              | 71         |
| I. T. Kurtsin, Studies in the Field of Viscerocortical Pathology                                                               | 77         |
| V. V. Parin, Trophic Guarantee of Cardiac Activity                                                                             | 89         |
| V. S. Rusinov and V. L. Ezorkhi, Prolonged Changes in the Membrane Potential of the Neurons and the Possible Functional Role   | 95         |
| A. D. Slonin, New Data on Physiological Adaptations to an Arid Zone                                                            | 101        |
| A. V. Solov'yev and L. I. Dvinyaninov, Pathogenesis of Experimental Gastric and Duodenal Ulcers in Dogs                        | 109        |
| New Data on Interoception Studies                                                                                              |            |
| M. G. Amiragova, Cerebral Cortex and Thyroid Gland                                                                             | 115        |
| N. A. Anikina, Resistance of Various Reactive Receptor Systems of the Small Intestines to Metabolic Disturbances               | 124        |

2/5

USSR

KREPS, Ye. M., Mekhanizmy regulyatsii fiziologicheskikh funktsiy, Sbornik statey k 40-letiyu nauchnoy deyatel'nosti akademika Vladimira Nikolayevicha Chernigovskogo, Nauka Press, Leningrad, 1971, 316 pp

|                                                                                                                                                            |             |
|------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|
| N. N. Beller, Mechanism of Efferent Effects of the Limbic Cortex on Autonomic Functions                                                                    | Page<br>132 |
| Ye. B. Berkhin, Introceptive Connections of the Kidneys                                                                                                    | 144         |
| R. A. Durinyan, Functional Significance of the Spatial Organization of the Visceral Afferent Systems on the Thalamocortical Level                          | 152         |
| V. Yu. Yermolayeva, Functional Organization of the Cortical and Corticofugal Paths of First and Second Somatosensory Zones of the Brain of a Cat           | 160         |
| O. B. Il'inskiy, Physiology of Single Mechanoreceptors (Pacinian Corpuscles)                                                                               | 173         |
| G. S. Kan, Study of the Role of the Introceptive Analyzer in the Pathogenesis of Certain Disturbances of Blood Circulation in the Presence of Tuberculosis | 185         |
| Ye. L. Kan, Study of the Mechanism of Participation of the Peripheral Innervation of the Spleen in Regulating the Erythrocyte Equilibrium                  | 192         |
| 3/5                                                                                                                                                        |             |



USSR

KREPS, Ye. M., Mekhanizmy regulyatsii fiziologicheskikh funktsiy, Sbornik statey k 40-letiyu nauchnoy deyatel'nosti akademika Vladimira Nikolayevicha Chernigovskogo, Nauka Press, Leningrad, 1971, 316 pp

|                                                                                                                                                                                                                                 |             |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|
| B. S. Kulayev, Significance of the Interoceptors in the Evolution of Vertebrates and Their Vital Activity                                                                                                                       | Page<br>203 |
| K. M. Kullanda, New Data on Analysis of the Properties and Mechanisms of the Genesis and Functional Significance of the Basic Forms of Evoked Potentials in the Somatovisceral Afferent Projection Zones of the Cerebral Cortex | 214         |
| T. S. Lagutina, Interaction of the Intero and Exteroceptors in the Process of Establishing the Reflex Acts of Urination and Defecation in Ontogenesis                                                                           | 228         |
| O. S. Merkulova, New Experimental Data With Respect to the Morphophysiological Analysis of the Adaptation Phenomenon in the Somatic Reflex Arc                                                                                  | 241         |
| S. S. Musyashchikova, Afferent Visceral Representation in the Central Nervous System                                                                                                                                            | 258         |
| T. V. Popova, Bioelectric Activity of the Medulla Oblongata in the Presence of Hypothermia and Blood Letting                                                                                                                    | 277         |

4/5

USSR

KREPS, Ye. M., Mekhanizmy regulyatsii fiziologicheskikh funktsiy, Sbornik statey k 40-letiyu nauchnoy deyatel'nosti akademika Vladimira Nikolayevicha Chernigovskogo, Nauka Press, Leningrad, 1971, 316 pp

|                                                                                                     | Page |
|-----------------------------------------------------------------------------------------------------|------|
| V. M. Khayutin, Level of Completion of Vasomotor Reflexes                                           | 292  |
| A. Ya. Yaroshevskiy, Erythropoiesis and Its Regulation in the Presence of Certain Internal Diseases | 302  |

5/5

81

USSR


UDC 547.26'118 + 547.341 + 547.345

NESTEROV, L. V., KREPYSHEYEVA, N. Ye., SABIROVA, R. A., and ROMANOVA, G. N.,  
Institute of Organic and Physical Chemistry imeni A. Ye. Arbutov, USSR Academy  
of Sciences

"Derivatives of Phosphorous Acid. VIII. Reaction of the Dialkyl Trialkyl-  
silyl Phosphites With Aldehydes and Ketones"

Leningrad, Zhurnal Obshchey Khimii, Vol XLI, No 11, 1971, pp 2,449-2,452

Abstract: Dialkyl acyl and tetraalkyl pyrophosphites are known to react with  
carbonyl compounds according to  $(RO)_2POAc + R'R''CO \rightarrow (RO)_2P(O)C(OAc)R'R''$   
(R = alkyl, R' = alkyl, aryl, R'' = H, alkyl and Ac = acyl or  $P(OR)_2$ ). In  
this connection, the reactions of dialkyl trialkylsilyl phosphites with  
aldehydes and ketones were studied. These reactions were found to proceed  
along exactly the same lines when Ac =  $SiAlk_3$ , and the reactions proceed under  
much less rigorous conditions when Ac = acyl. The aldehydes react rapidly,  
with release of heat, while the ketones require preheating to 100°C. In all  
cases, there is formation of the dialkyl esters of  $\alpha$ -(trialkylsiloxy)alkyl-  
phosphonic acids:



1/2

USSR

NESTEROV, L. V., et al., Zhurnal Obshchey Khimii, Vol XLI, No 11, Nov 1971, pp 2,449-2,452

These compounds have been known only for a few years, and until now no satisfactory method had been devised for obtaining them. Physical data for the nine compounds obtained, along with laboratory procedures, are given in the paper.

2/2

- 29 -

USSR

UDC 546.691.632.653:543.757.1

KRESHKOV, A. P., SAYUSHKINA, Ye. N., and TIMASHEVA, N. A.,  
Moscow Institute of Chemical Technology named D. I. Mendeleev  
"Analysis of Binary Mixtures of Gallium, Indium, and Thallium  
Salts by Methods of Potentiometric Titration and Ion Exchange  
in Non-Aqueous Solutions"

Moscow, Zavodskaya Laboratoriya, Vol 37, No 12, 1971,  
pp 1421-1424

Abstract : The behavior of chlorides, nitrates, and sulfates of  
gallium (III) and indium (III), nitrate of thallium (I), and  
accompanying elements [Zn (II), Cd (II), Cu (II), Co (II)] in  
non-aqueous solutions was investigated. For the quantitative  
determination of nitrates and thallium and indium chlorides  
and their accompanying elements, a mixed solvent, water-aceto-  
nic (1:4), was selected in which the salts show like acids and  
are titrated potentiometrically by 0.1 ethanolic solution of

1/2

USSR

KRESHKOV, A. P., et al., Zavodskaya Laboratoriya, Vol 37, No 12, 1971,  
pp 1421-1424

KOH. Thallium nitrate develops a neutral character in water-acetone media. For its determination, the ion exchange on cationite SVD-3 in H-form is used. For the determination of telluric and indium sulfates and other elements and also mixtures of sulfates with nitrates ( or chlorides ), the cationite SVD-3 in H-form is used with following potentiometric titration of generated acids by 0.1-normal methanol solution of diphenylguanidine. Determinations of salts and their mixtures are described. Two illustrations, two tables, 13 biblio. refs.

2/2

- 65 -

1/2 015 UNCLASSIFIED PROCESSING DATE--04DEC70  
TITLE--DIFFERENTIATED DETERMINATION OF XYLENE AND TOLUENE ISOMERS IN  
NONAQUEOUS SOLVENTS -U-  
AUTHOR-(03)-KRESHKOV, A.P., GURETSKIY, I.YA., SHOLOVA, N.T.  
COUNTRY OF INFO--USSR  
SOURCE--MATER. IKH PRIMEN. 1970, (2), 53-6  
DATE PUBLISHED-----70  
  
SUBJECT AREAS--CHEMISTRY  
TOPIC TAGS--XYLENE, TOLUENE, OXIDATION, CARBOXYLIC ACID, POTENTIOMETRIC  
TITRATION, ISOMER  
  
CONTROL MARKING--NO RESTRICTIONS  
DOCUMENT CLASS--UNCLASSIFIED  
PROXY FICHE NO----FD70/605012/D08 STEP NO--UR/0303/70/000/002/0053/0056  
CIRC ACCESSION NO--AP0140297

UNCLASSIFIED

2/2 015

UNCLASSIFIED

PROCESSING DATE--04DEC70

CIRC ACCESSION NO--AP0140297

ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. MIXTS. OF PMME AND O, M, AND P XYLENE WERE OXIDIZED 40-5 MIN WITH KMNO SUB4 SOLN. AT 95-80DEGREES TO GIVE BZOH AND THE CORRESPONDING DIBASIC ACIDS. IR SPECTRA AND GAS CHROMATOG. SHOWED THAT ONLY CARBOXYLIC ACIDS WER FORMED, THERE WAS NO DECARBOXYLATION. EACH PRODUCT GIVES A CHARACTERISTIC POTENTIOMETRIC TITRN. CURVE AND CAN BE DETD. IN THE PRESENCE OF OTHER COMPONENTS. THE TITRNS. WERE CARRIED OUT IN AN ACETONE WATER MECH ISO PROH MIXT. AGAINST ISO PROK. THE MEAN ARITHMETIC DETN. ERROR WAS LESS THAN OR EQUAL TO 5.36PERCENT.

UNCLASSIFIED



Acc. Nr.

AP0049838

Abstracting Service:  
CHEMICAL ABST. 5/70

Ref. Code

UR 0075

K

106738y Theoretical curves of the conductometric titration of acids and bases in nonaqueous solutions. Khudyakova, T. A.; Kreshkov, A. P. (Gorki Polytech. Inst., Gorki, USSR). Zh.

Anal. Khim. 1970, 25(1), 11-17 (Russ). Equations were derived for titrn. curves of acids and bases in nonaq. solvents, permitting the calcn. of equil. concns. of solvated H ions, acid anions, and base cations during the titrn. Ion concn. dependences were found and theoretical curves of the conductometric titrn. of acids of different strength by strong and weak bases in a nonaq. solvent with  $pK_a = 20$  were derived. Chaim Weiner

1/1

REEL/FRAME  
19801763

714

Acc. Nr. **AP0053426** Abstracting Service:  
CHEMICAL ABST.

Ref. Code:

110619c Acid-base properties of alkyl(aryl)fluorosilanes in nonaqueous media. Drozdov, V. A.; Kreshkov, A. B.; Rumanova, A. D. (USSR). *Zh. Obshch. Khim.* 1970, 40(1), 104-9 (Russ). Conventional syntheses gave the following:  $\text{Me}_3\text{HSiF}$ , b.  $51^\circ$ , d $^{20}_4$  —,  $n_D^{20}$  — ( $n_D^{20}$  1.3570);  $\text{Me}_2\text{PrSiF}$ , b.  $77^\circ$  —, 1.3708;  $\text{Me}_2\text{PhSiF}$ , b.  $155-7^\circ$  —, 1.5471;  $(\text{PhCH}_2)_2\text{SiMe}_2\text{F}$ , b.  $104^\circ$ , 0.9743, 1.4838;  $\text{MeEt}(\text{PhCH}_2)_2\text{SiF}$ , b.  $200-2^\circ$ , 0.9069, 1.4702;  $\text{MePh}(\text{PhCH}_2)_2\text{SiF}$ , b.  $117^\circ$ , 1.0610, 1.5481;  $\text{MePh}_2\text{SiF}$ , b.  $164.5^\circ$ , 1.077, 1.5482;  $\text{Et}_2\text{SiF}$ , b.  $109-10^\circ$ , 0.8303, 1.3906;  $\text{Et}_2\text{PhSiF}$ , b.  $77-8^\circ$ , 0.9837, 1.4880;  $\text{Et}(\text{PhCH}_2)_2\text{SiF}$ , b.  $90^\circ$ , 0.9828, 1.4942;  $\text{PhSiFCl}_2$ , b.  $157-9^\circ$  —, 1.5203;  $\text{Ph}_2\text{SiFCl}$ , b.  $131-2^\circ$  —, 1.5471;  $\text{MePhSiF}_2$ , b.  $108-10^\circ$ , 1.2000, 1.5339;  $\text{PhSiF}_2\text{Cl}$ , b.  $129-30^\circ$  —, 1.4575;  $\text{PhSiF}_3$ , b.  $102-5^\circ$ , 1.2270, 1.4106; and  $\text{PhSiCl}_3$ , b.  $225^\circ$ , 1.2220, 1.5821; also:  $\text{MePh}_2\text{SiCl}$ , b.  $191^\circ$ ;  $\text{MePhSiCl}_2$ , b.  $223-5^\circ$ ;  $\text{MePh}(\text{PhCH}_2)_2\text{SiCl}$ , b.  $149^\circ$ ;  $\text{Me}_2\text{PhSiCl}$ , b.  $190-1^\circ$ ; and  $\text{Et}_2\text{PhSiCl}$ , b.  $100-2^\circ$ . Ir spectral curves of 12 of these were shown and their thermodynamic acidity was detd. by potentiometric titrn. in dry MeOH and EtOH relative to dry HCl in the same solvents. The relative values of such acidity expressed in units of pK (— $\text{H}^+\text{R}_1$ ) were, in MeOH and EtOH, resp.:  $\text{R}_2\text{SiF}$ , 10.0, 10.0;  $\text{MeEt}(\text{PhCH}_2)_2\text{SiF}$ , 10.2, 9.50;  $\text{Me}_2\text{PhSiF}$ , 9.55, 9.14;  $\text{MePh}(\text{PhCH}_2)_2\text{SiF}$ , 9.29, 8.80; and  $\text{Me}_2\text{SiF}$ , 10.43, 9.69. Unlike chlorosilanes, the substituent groups had considerable effect on the acidity of the fluorosilanes. This is ascribed to the low polarizability of these.

G. M. Kizilevskii

REEL/FRAME  
19830451

1/2 010 UNCLASSIFIED PROCESSING DATE--16OCT70  
TITLE--CHEMICAL ANALYTICAL BEHAVIOR OF SULFUR CONTAINING ALIPHATIC  
DICARBOXYLIC ACIDS IN NONAQUEOUS SOLVENTS -U-  
AUTHOR-(03)-KRESHKOV, A.P., ALGAROVA, N.SH., TANGANDY, B.B.  
COUNTRY OF INFO--USSR  
SOURCE--ZH. ANAL. KHIM. 1970, 25(2), 362-8  
DATE PUBLISHED-----70  
SUBJECT AREAS--CHEMISTRY  
TOPIC TAGS--ORGANIC SOLVENT, ALIPHATIC DICARBOXYLIC ACID, CHEMICAL  
ANALYSIS, THIOL, DISSOCIATION CONSTANT, POTENTIOMETRIC TITRATION  
CONTROL MARKING--NO RESTRICTIONS  
DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FRA--1993/0685 STEP NO--UR/0075/70/025/002/0362/0365  
CIRC ACCESSION NO--AP0113956  
UNCLASSIFIED

2/2 010

UNCLASSIFIED

PROCESSING DATE--16OCT70

CIRC ACCESSION NO--AP0113556

ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. POTENTIOMETRIC TITRN. OF S CONTG. DICARBOXYLIC ACIDS IN A MEQH, ETOH, HEXANOL, ACETOPHENONE, HCONME SUB2, AND MECH WAS STUDIED. ACIDITY CONSTS., PK SUBA FOR BETA, BETA PRIME, THIODIPROIONIC, DELTA, DELTA PRIME, THIODIYALERIC, OMEGA, OMEGA PRIME, THIODIENANTHIC, OMEGA, OMEGA PRIME, THIODIPELARGONIC, AND OMEGA, OMEGA PRIME, THIODIUNDECANIC ACIDS IN THE ABOVE SOLVENTS WERE EVALUATED. THE CONDITIONS FOR THE DETN. OF OMEGA, OMEGA PRIME, THIODICARBOXYLIC ACIDS IN NONAQ. SOLVENTS BY THE CALCN. OF THE TITRN. CONST., PK SUBT, VALUES WERE EXPLAINED. ALIPHATIC MONOHYDROIC NORMAL ALCS. HAVE A LEVELING ACTION ON OMEGA, OMEGA PRIME, THIODICARBOXYLIC ACIDS, WHICH BEHAVE IN ALC. SOLNS. AS WEAK ACIDS (PK SUBA EQUALS 9-11). THE DEPENDENCES PK SUBA-PK SUBS (CONST. OF SOLVENT AUTOPROTOLYSIS) AND PK SUBT-PK SUBS FOR THE STUDIED COMPS. HAVE A LINEAR CHARACTER, PROVING THE EFFECT OF THE SOLVENT ON THE ACID BASE PROPERTIES AND CONDITIONS OF QUANT. DETN. OF THE ACIDS. THE SEP. DETN. OF THE 2 CARBOXYL GROUPS OF DICARBOXYLIC ACIDS CONTG. S IN THE C CHAIN INCREASE IN THE SERIES OF HCONME SUB2 IS SMALLER THAN MECH IS SMALLER THAN ME SUB2 CO. A METHOD WAS DEVELOPED FOR THE QUANT. DETN. OF DICARBOXYLIC ACIDS IN DIFFERENTIATING SOLVENTS WITH A STD. C SUB6 H SUB6-MEQH SOLN. OF ET SUB4 NOH. FACILITY: MOSCOW CHEM. TECHNOL. INST., MOSCOW, USSR.

UNCLASSIFIED

AA0044227

K

UR 0482

Soviet Inventions Illustrated, Section I Chemical, Derwent, 3-70

228316 DETERMINATION OF BOUND SULPHURIC ACID in  
cellulose esters during their prod. can be  
carried out by a rapid, accurate and highly sensit-  
ive method consisting of dissolving a sample of the  
material to be analysed in a chloroform-methanol  
mixture, adding a solution of methylene blue in  
aq. HCl, and photometrically analysing the coloured  
organic phase, which separates. In an example, a  
100 mg sample was dissolved in a 10:1 chloroform  
methanol mixture in a 100 ml. measuring flask;  
a 10 ml. aliquot portion was mixed with 10 ml. of  
the 10:1 chloroform-methanol mixt. and 40 ml. of a  
soln. of methylene blue (10 mg/l) in 0.01 N HCl;  
the mixture was shaken for 10 secs, and after 10  
mins the organic phase was separated and subjected  
to photometric analysis. 24.7.67. as 1174844/21-4  
KRESHKOV A P. et al. Vladimir Synthetic Resins  
Res. Inst (25.6.67) Bul. 31/8.10.68. Class 421  
Int. Cl. G 01a. 1

19770721

AA0044227

Kreshkov, A. P.; Yemelin, Ye. A.; Yavgrafova, I. V.; Solov'yeva, O. A.

Vladimirskiy Nauchno - Issledovatel'skiy

Institut Sinteticheskikh Smol

19770722

1/2 009 UNCLASSIFIED PROCESSING DATE--27NOV70  
TITLE--EVALUATION OF NONAQUEOUS SOLVENT AUTOPROTOLYSIS CONSTANTS -U-  
AUTHOR--(04)-KRESHKOV, A.P., ALDAROVA, N.SH., SMOLOVA, N.T., RYAGUZOV, A.I.  
COUNTRY OF INFO--USSR  
SOURCE--ZH. FIZ. KHIM. 1970, 44(4), 1126-7  
DATE PUBLISHED-----70  
SUBJECT AREAS--CHEMISTRY  
TOPIC TAGS--ALIPHATIC ALCOHOL, GLYCOL, KETONE, DICARBOXYLIC ACID, SOLVENT  
ACTION  
CONTROL MARKING--NO RESTRICTIONS  
DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FRAME--3007/0795 STEP NO--UR/0076/70/064/004/1126/1127  
CIRC ACCESSION NO--AP0136229  
UNCLASSIFIED

UNCLASSIFIED

PROCESSING DATE--27NOV70

2/2 009

CIRC ACCESSION NO--AP0136229

ABSTRACT/EXTRACT--(U) GP-O-

ABSTRACT. THE AUTOPROTOLYSIS CONSTS., PK  
SUBS, WERE ESTD. FOR A NO. OF ALIPHATIC ALCS. (MEOH TO OCTANOL),  
GLYCOLS, AND KETONES FROM THE LINEAR RELATIONSHIP PK SUBS VS. PA SUBS,  
WHERE PK SUBA HAS THE ACID CONST. OF A DICARBOXYLIC ACID IN A GIVEN  
SOLVENT GROUP, AND FROM THE EXPRESSION PK SUBS EQUALS E SUBS -K SUBS,  
WHERE E SUBS IS THE RELATIVE ACIDITY SCALE AND K SUBS THE COEFF. OF THE  
RELATIVE ACIDITY SCALE. THE K SUBS VALUES FOR THE SOLVENTS DECREASED  
IN THE ORDER KETONES GREATER THAN ALCS. GREATER THAN GLYCOLS GREATER  
THAN H SUB2 O.  
MENDELEEVA, MOSCOW, USSR.

FACILITY: MOSK. KHIM.-TEKHNO. INST. IM.

UNCLASSIFIED



1/2 007 UNCLASSIFIED PROCESSING DATE--30OCT70  
TITLE--EVALUATION OF AUTOPROTOLYSIS CONSTANTS FOR NONAQUEOUS SOLVENTS BY  
CORRELATION EQUATIONS, PK, F PKA MH -U-  
AUTHOR--(04)--KRESHKOV, A.P., ALDAROVA, N.SH., TANGANOV, B.B.,  
SLAVGRODSKAYA, N.V.  
COUNTRY OF INFO--USSR  
SOURCE--ZH. FIZ. KHIM. 1970, 44(1), 241-3  
DATE PUBLISHED--70

SUBJECT AREAS--CHEMISTRY

TOPIC TAGS--HYDROGEN ION CONCENTRATION, PROPANOL, BUTANOL, ACETONE,  
SOLVENT ACTION

CONTROL MARKING--NO RESTRICTIONS

DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FRA--2000/0875

STEP NO--UR/0076/70/044/001/0241/0243

CIRC ACCESSION NO--AP0124538

UNCLASSIFIED

2/2 007

UNCLASSIFIED

PROCESSING DATE--30OCT70

CIRC ACCESSION NO--AP0124538

ABSTRACT/EXTRACT--(U) GP-O- ABSTRACT. THEORETICAL RELATIVE ACIDITIES WERE CALCD. FOR NON AQ. SOLVENTS. A LINEAR CORRELATION PK SUBA EQUALS 0.0145, E SUBB 9.71 (E SUBB IS THE RELATIVE ACIDITY) WAS FOUND FOR ALKANOLS. AUTOPROTOLYSIS CONSTS. ARE LINEARLY RELATED TO PK SUBA; PK SUBS EQUALS 0.843, PK SUBA 1.74. THIS EQUATION WAS USED FOR THE PREDICTION OF THEORETICAL ACIDITIES OF PROH, BUCH, N,HEXANOL, N,HEPTANOL, N,OCTANOL, ACETONE, MECOET, AND MECOPR. THE AUTOPROTOLYSIS CONSTS. OF THE COMPS. NOT USED IN THE CORRELATION ARE IN GOOD AGREEMENT WITH PUBLISHED DATA. FACILITY: MOSK. KHIM. TEKHNOL. INST. IM. MENDELEEVA, MOSCOW, USSR.

UNCLASSIFIED

1/2 017 UNCLASSIFIED PROCESSING DATE--30OCT70  
TITLE--EFFECT OF ACETONITRILE AND WATER ON THE DIFFERENTIATING ACTION OF  
ACETONE WITH RESPECT TO PHTHALIC ACID ISOMERS -U-  
AUTHOR--(04)--KRESHKOV, A.P., GURETSKIY, I.YA., SMOLOVA, N.T., RYAGUZOV,  
A.I.

COUNTRY OF INFO--USSR

SOURCE--ZH. ANAL. KHIM. 1970, 25(3), 451-7

DATE PUBLISHED--70

SUBJECT AREAS--CHEMISTRY

TOPIC TAGS--ACETONITRILE, ACETONE, PHTHALIC ACID, ISOMER, CALCULATION,  
ELECTROLYTE, POTENTIOMETRIC TITRATION

CONTROL MARKING--NO RESTRICTIONS

DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FRA--2000/2036

STEP NO--UR/0078770/025/003/0451/0457

CIRC ACCESSION NO--AP0125443

UNCLASSIFIED

2/2 017

UNCLASSIFIED

PROCESSING DATE--000CT70

CIRC ACCESSION NO--AP0125643

ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. AN EQUATION FOR CALC. THE PK OF ELECTROLYTES IS SUGGESTED WHICH CAN BE APPLIED ONLY IF THE POTENTIOMETRIC TITRN. IS ACCOMPANIED BY PPT. FORMATION. BY MEANS OF THIS EQUATION, THE PK OF O, M, AND P, PHTHALIC ACIDS WERE CALCD. THE EFFECTS OF H SUB2 O AND MECH ON THE DIFFERENTIATING ACTION OF ME SUB2 CO WITH RESPECT TO PHTHALIC ACID ISOMERS WERE STUDIED. H SUB2 DECREASES THE ACTION AND MECH HAS NO ESSENTIAL EFFECT ON THE DIFFERENTIATING ACTION OF ME SUB2 CO. THIS WAS TAKEN INTO ACCOUNT DURING THE DEVELOPMENT OF THE METHOD FOR ANALYZING MIXTS. OF PHTHALIC ACID ISOMERS IN A MIXED ME SUB2 CO, MECH, H SUB2 O SOLVENT. FACILITY: MOSCOW CHEM.-TECHNOL. INST., MOSCOW, USSR.

UNCLASSIFIED

US3R

UDC: 621.396.6-181.5

KRESIN, O. M., STAROS, F. G., YAKOVLEV, A. S.

"A Method of Determining the Temperature Fields of Planar Systems"

Elektron. tekhnika. Nauchn.-tekhn. sb., Mikroelektronika (Electronic Technology. Scientific and Technical Collection. Microelectronics), 1970, vyp. 5(26), pp 105-113 (from RZh-Radiotekhnika, No 5, May 71, Abstract No 5V216)

Translation: Methods are developed for calculating and analyzing the temperature fields of fuel elements and groups of elements of arbitrary topology. The proposed method is applicable to calculations in thin-film and integrated microcircuits and mosaic circuit assemblies, as well as in designs which utilize ferrite films. Resumé.

1/2 035 UNCLASSIFIED PROCESSING DATE--04DEC70  
TITLE--BEHAVIOR OF PLUTONIUM-239 IN RABBITS DURING THE INHALATION OF  
AMMONIUM PENTACARBONATOPLUTONATE -U-  
AUTHOR--(02)-TSEVELEVA, I.A., KRESLOV, V.V.  
COUNTRY OF INFO--USSR  
SOURCE--RADIOBIOLOGIYA 1970, 10(2), 307  
DATE PUBLISHED-----70  
SUBJECT AREAS--BIOLOGICAL AND MEDICAL SCIENCES  
TOPIC TAGS--PLUTONIUM ISOTOPE, RABBIT, HALF LIFE, LUNG, BLOOD CHEMISTRY,  
LIVER, BONE, RADIATION DOSAGE  
CONTROL MARKING--NO RESTRICTIONS  
DOCUMENT CLASS--UNCLASSIFIED  
PROXY FICHE NO----FD70/605004/811 STEP NO--UR70205/70/010/002/0307/0307  
CIRC ACCESSION NO--AP0139607  
UNCLASSIFIED

2/2 035

UNCLASSIFIED

PROCESSING DATE--04DEC70

CIRC ACCESSION NO--AP0139607

ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. DISTRIBUTION AND KINETICS OF PRIME239 PU FOLLOWING A SINGLE INHALATION OF (INH SUB4) SUB5 PU (CO SUB3) SUB5 (1.5 AND 5.0 MU CI) WAS STUDIED IN RABBITS OVER 2 YEARS. REMOVAL OF PRIME239 PU FROM THE LUNGS DID NOT DEPEND SIGNIFICANTLY ON THE AMT. OF RADIOISOTOPE ADMINISTERED AND WAS DESCRIBED WITH THE HELP OF 3 EFFECTIVE HALF LIVES (T SUBEFF): 61.5PERCENT WITH T SUBEFF EQUALS 3 DAYS, 35.2PERCENT WITH T SUBEFF EQUALS 35 DAYS, AND 2.7PERCENT WITH T SUBEFF EQUALS 174 DAYS. ABOUT 10-15PERCENT OF THE DOSE ADMINISTERED WAS REMOVED FROM THE LUNGS INTO THE BLOOD AND WAS THEN DEPOSITED IN APPROX. EQUAL AMTS. IN THE LIVER AND SKELETON. THE CONTENT OF PRIME239 PU DID NOT CHANGE IN THESE ORGANS SIGNIFICANTLY DURING 2 YEARS. THE ABSORBED DOSE OF RADIATION IN THE LUNGS, LIVER AND SKELETON, CALCD. FOR THE PERIOD OF 2 YEARS, WAS 3240, 490, AND 165 RADS, RESP. FACILITY: INST. BIOFIZ., MOSCOW, USSR.

UNCLASSIFIED

USSR

UDC 539.4.012

KRESTIN, G. S., LIBATSKIY, L. L., YAREMA, S. YA., Physical Mechanics Institute  
of the Ukrainian SSR Academy of Sciences, L'vov, L'vov Polytechnic Institute

"Stressed State of a Disc with a Diametral Crack"

Fiziko-Khimicheskaya Mekhanika Materialov, Vol 8, No 2, 1972, pp 69-73

Abstract: A solution of the integral equation of the symmetric problem in the form of a series with respect to  $\lambda$  ( $\lambda$  is the crack length reduced to the disc diameter) is found for a disc with a central diametral crack with arbitrary symmetric and antisymmetric loads. Formulas are also found for determining the stress intensity coefficients with accuracy to terms of order  $\lambda^{1/2}$ . Graphs are presented showing the variation of the critical value of concentrated normal and tangential forces as a function of the crack length and the variation of the corresponding distributed loads.

1/1



USSR

UDC 542.91:547.1'118:541.69

KRESTKIN, A. P., VOLKOVA, R. I., GODOVIKOV, N. N., KABACHNIK, M. I.,  
KARDANOV, N. A., Institute of Organoelemental Compounds of the Soviet  
Academy of Sciences; Institute of Evolutionary Physiology and Biochemistry  
imeni I. M. Sechenov of the Soviet Academy of Sciences

"Interaction of S-( $\beta$ -Alkylmercaptoethyl)diphenylthiophosphinates,  
O-( $\beta$ -Alkylmercaptoethyl)diphenylphosphinates and Their Iodomethylates  
With Butyrylcholinesterase"

Moscow, Izv. An SSSR: Ser. Khimicheskaya, No 3, Mar 73, pp 622-627

Abstract: The paper gives the results of research on the kinetics of  
inhibition of butylcholinesterase by S-  $\beta$ -alkylmercaptoethyl diphenyl-  
thiophosphinates, O-  $\beta$ -alkylmercaptoethyl diphenylphosphinates and their  
methiodides. It is found that these substances have a combination type  
of inhibiting action which increases with the length of the alkyl radical  
bound to the sulfide or sulfonium sulfur atom.

1/1

- 21 -

USSR

UDC: 681.396.96:681.373

KRESINIKOV, L. A.

"Experimental Studies of Radar Characteristics of Hydrometeorological Formations in the Millimeter Range"

Tr. Sev.-Zap. nauch. politekhn. in-t (Works of the Northwest Polytechnical Correspondence Institute), 1972, No 20, pp 12-13 (from Tekhn. Radiotekhnika, No 12, Dec 72, abstract No 12625 [résumé])

Translation: The paper describes equipment for studying the radar characteristics of various objects by the method of hydroacoustic modeling, and gives the measurement procedure.

1/1

USSR

UDC 54-162.2:546.799.4

KRESTOV, G. A., and KRESTOVA, N. V.

"Temperature of Coefficients and Variations in the Interionic Distance of Crystalline Compounds of Plutonium at Various Temperatures"

Leningrad, Radiokhimiya, Vol 12, No 4, 1970, pp 600-606

Abstract: The authors investigated temperature coefficients and changes in interionic distances as a function of temperature for some plutonium halides and chalcogenides. The data necessary for the computations were taken from recent literature on this class of compounds. It was found that the numerical values of the ionic components of the temperature coefficients and changes in interionic distances of the investigated salts increase with a rise in temperature. This increase is comparatively small over a considerable temperature range. The maximum change in the radius of the trivalent plutonium ion is 0.02 Å, while that of the tetravalent plutonium ion is 0.01 Å. The radii of the halide and chalcogenide ions vary over the same range. The change in interionic distances from 293.15°K to the melting point of the salt reaches 0.05 Å.

1/1

1/2 018 UNCLASSIFIED PROCESSING DATE--09OCT70  
TITLE--CHANGES IN THE ENTROPY COMPONENT OF FREE ENERGY CHANGE DURING THE  
HYDRATION OF MONATOMIC IONS AT VARIOUS TEMPERATURES -U-  
AUTHOR--(02)-KRESTOV, G.A., ABRASIMOV, V.K.

COUNTRY OF INFO--USSR

SOURCE--IZV. VYSSH. UCHEB. ZAVED., KHIM. KHIM. TEKHNOL. 1970, 13(1), 41-3

DATE PUBLISHED-----70

SUBJECT AREAS--CHEMISTRY

TOPIC TAGS--HYDRATION, ENTROPY, WATER, THERMAL EFFECT, MOLECULAR STRUCTURE

CONTROL MARKING--NO RESTRICTIONS

DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FRA--1993/1919

STEP NO--08/0153/70/013/001/0041/0043

CIRC ACCESSION NO--AT0114359

UNCLASSIFIED

2/2 018

UNCLASSIFIED

PROCESSING DATE--09OCT70

CIRC ACCESSION NO--AT0114359

ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. A TABLE IS PRESENTED OF THE CHANGES IN THE ENTROPY COMPONENTS OF THE FREE ENERGY CHANGE,  $T \Delta S_{SUBI}$ ,  $T \Delta S_{SUBII}$ ,  $T \Delta S_{SUB(LONG RANGE)}$ , AND  $T \Delta S_{SUB(SHORT RANGE)}$ , FOR HYDRATION OF THE IONS LI POSITIVE, NA POSITIVE, K POSITIVE, RB POSITIVE, CS POSITIVE, CA PRIME2POSITIVE, SI PRIME2POSITIVE, BA PRIME2POSITIVE, CD PRIME2POSITIVE, CL NEGATIVE, BR NEGATIVE, I NEGATIVE AT 0, 25.15, 40, 60, 80, AND 100DEGREES. THE BASIC INFLUENCE ON THE TEMP. DEPENDENCE OF THE ENTROPY OF HYDRATION IS THAT OF THE STRUCTURAL COMPONENT  $T \Delta S_{SUBII}$  AND PARTICULARLY THAT OF  $T \Delta S_{SUB(SHORT RANGE)}$ . STRUCTURAL CHANGES OF WATER WITH INCREASING TEMP. INHIBIT IONIC HYDRATION. FACILITY: IVANDV. KHIM.--TEKHNOL. INST., IVANOVO, USSR.

UNCLASSIFIED

2/2 026

UNCLASSIFIED

PROCESSING DATE--30OCT70

CIRC ACCESSION NO--AT0116980

ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. THE OSTWALD SOLY. COEFF. 10 PRIME3  
GAMMA OF AR IN H SUB2 O DECREASES FROM 30 AT 25DEGREES TO 19 AT  
70DEGREES. FOR SOLNS. CONTG. 0.25 MOLE FRACTION (CH SUB2 OH) SUB2 IN H  
SUB2 O, 10 PRIME3 GAMMA IS 27 ESSENTIALLY INDEPENDENT OF TEMP. AT  
HIGHER CONCNS. OF GLYCOL, THE SOLY. OF AR INCREASES WITH TEMP., I.E.,  
FOR 0.6 MOLE FRACTION GLYCOL FROM 10 PRIME3 GAMMA EQUALS 28 AT 25DEGREES  
TO 37 AT 70DEGREES. THERMODYNAMIC DATA ARE CALCD. FROM THE SOLY. TEMP.  
RELATIONSHIPS. DELTA F SUBSOLN. SHOWS LITTLE VARIATION WITH CONCEN. OF  
(CH SUB2 OH) SUB2 AND IS SIMILAR TO 4000 CAL-MOLE AT 25DEGREES AND 4700  
CAL-MOLE AT 70DEGREES. DELTA S DEGREES AT 70DEGREES DECREASES FROM 5700  
CAL-MOLE IN H SUB2 O TO 3500 CAL-MOLE IN H SUB2 O CONTG. 0.7 MOLE  
FRACTION GLYCOL. ABOVE A 0.15 MOLE FRACTION GLYCOL, DELTA H DEGREES  
SUBSOLN. SHOWS LITTLE EFFECT OF TEMP. AND INCREASES FROM MINUS 500  
CAL-MOLE AT 0.15 MOLE FRACTION TO 1000 CAL-MOLE AT 0.7 MOLE FRACTION.  
IN THE SAME RANGE, DELTA S DEGREES SUBSOLN. INCREASES FROM MINUS 15 TO  
MINUS 10 ENTROPY UNITS. FACILITY: IVANOV. KHIM.-TEKHNOL. INST.,  
IVANOVO, USSR.

UNCLASSIFIED

Acc. Nr.

AP0034105

Abstracting Service:  
CHEMICAL ABST. 4-70

Ref. Code

NR 0078

74180s Derivatographic method for determining the coordination numbers of some trivalent ions of rare earth elements. A. A. Kurnikina, G. I. (USSR). Zh. Neorg. Khim. 1970, 15(1), 73-6 (Russ). A new method was developed for detn. of coordination nos. of ions of rare earth elements by DTA. Tri-valent Pr, Nd, Sm, Eu, Gd, Er, Yb, and Y ions are 6 coordinated. Calcn. of entropy (S) of a mole of H<sub>2</sub>O close to a 3-valent ion, and of a mole of H<sub>2</sub>O in pure solvent (S<sub>0</sub> and S<sub>1</sub>, resp.) showed that, with increasing radius of cations the value of S<sub>0</sub> decreases and approaches the value of S of ice (9.4 e.u.).

HMJR

REEL/FRAME

19710748

USSR

UDC 54-162.2:546.759.4

KRESTOV, G. A., and KRESTOVA, N. V.

"Temperature of Coefficients and Variations in the Interionic Distance of Crystalline Compounds of Plutonium at Various Temperatures"

Leningrad, Radiokhimiya, Vol 12, No 4, 1970, pp 600-605

Abstract: The authors investigated temperature coefficients and changes in interionic distances as a function of temperature for some plutonium halides and chalcogenides. The data necessary for the computations were taken from recent literature on this class of compounds. It was found that the numerical values of the ionic components of the temperature coefficients and changes in interionic distances of the investigated salts increase with a rise in temperature. This increase is comparatively small over a considerable temperature range. The maximum change in the radius of the trivalent plutonium ion is 0.02 Å, while that of the tetravalent plutonium ion is 0.01 Å. The radii of the halide and chalcogenide ions vary over the same range. The change in interionic distances from 298.15°K to the melting point of the salt reaches 0.05 Å.

1/1



USSR

UDC 534.1+539.3

BYKOVTSSEV, G. I., (Voronezh), and KRETOVA, L. D., (Voronezh)

"Shock Waves Propagation in Elastic Plastic Media"

Moscow, Prikladnaya Matematika i Mekhanika, Vol 36, No 1, 1972, pp 106-116

Abstract: The problem of shock wave propagation in elastic-plastic media is considered. It is shown that neutral shock waves on which plastic deformations are continuous, and waves on which plastic deformations are discontinuous can exist in ideal and hardening elastic media. Conditions of existence of the second type waves are formulated and velocities of all mentioned waves in ideal-plastic bodies are determined at arbitrary prominent yield and Tresca conditions, and also in hardening bodies at kinematic and isotropic hardening. Basic relationships for discontinuities in transition through waves surfaces are derived. The behavior of shock waves in propagation process under the von Mises and Tresca yield conditions is investigated using the second order kinematic compatibility conditions. It is shown that shock waves intensities vary in accordance with laws of geometric optics.

1/1

1/2 011 UNCLASSIFIED PROCESSING DATE--0900170  
TITLE--MOLECULAR WEIGHT OF MALATE DEHYDROGENASE OF SOYBEAN COTYLEDONS -U-  
AUTHOR-(02)-BRONJVITSKAYA, Z.S., KRETOVICH, Y.I. K  
COUNTRY OF INFO--USSR  
SOURCE--DOKL. AKAD. NAUK SSSR 1970, 190(2), 461-3  
DATE PUBLISHED-----70

SUBJECT AREAS--BIOLOGICAL AND MEDICAL SCIENCES  
TOPIC TAGS--AGRICULTURE CROP, DEHYDROGENASE, ULTRACENTRIFUGE, GEL,  
FILTRATION

CONTROL MARKING--NO RESTRICTIONS

DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FRA--1988/0936

STEP NO--UR/0020/70/190/002/0461/0463

CIRC ACCESSION NO--AT0105805

UNCLASSIFIED

2/2 011

UNCLASSIFIED

PROCESSING DATE--09OCT70

CIRC ACCESSION NO--AT0105805

ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. ULTRACENTRIFUGATION OF THE TITLE MATERIAL YIELDED 2 FORMS OF THE ENZYME FROM THE LEAF MATERIAL, ONE FRACTION HAVING A MOL. WT. OF SIMILAR TO 275,000, THE OTHER 69,000. ADJUSTMENT OF PH TO 6.2 OR TREATMENT WITH CSCL, NACL, OR UREA DISSOC. THE FORMER FRACTION INTO THE LATTER FORM. AT A LOW CONCN. OF NACL (0.1N) THIS DID NOT OCCUR. USING THE TAGGED PROTEIN TECHNIQUE AND GEL FILTRATION THE SPECIMENS OF THE TITLE ENZYME DERIVED FROM THE COTYLEDONS WAS FOUND TO HAVE MOL. WTS. OF SIMILAR TO 140,000-145,000 IN THE HIGHER FRACTION AND 67,000-70,000 IN THE LOWER FRACTION. EVIDENTLY THESE ARE TETRAMER AND DIMER UNITS. FACILITY: INST. BIOKHIM. IM. BAKHA, MOSCOW, USSR.

UNCLASSIFIED

1/2 - 010 UNCLASSIFIED PROCESSING DATE--090CT70  
TITLE--INFLUENCE OF LIGHT, DARKNESS AND NITROGEN NUTRITION ON KETO AND  
AMINOACIDS CONTENTS IN KIDNEY BEAN SPROUTS -U-  
AUTHOR--(04)--KRETOVICH, V.L., GEYKO, N.S., TSENOVA, M.P., VERKHOTURJOVA,  
G.S.  
COUNTRY OF INFO--USSR  
SOURCE--IZVESTIYA AKADEMII NAUK SSSR, SERIYA BIOLOGICHESKAYA, 1970, NR 2,  
PP 302-305  
DATE PUBLISHED-----70

SUBJECT AREAS--AGRICULTURE, BIOLOGICAL AND MEDICAL SCIENCES

TOPIC TAGS--PLANT PHYSIOLOGY, LEGUME CROP, AMINO ACID, NITROGEN, NUTRITION

CONTROL MARKING--NO RESTRICTIONS

DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FRA--1996/0475

STEP NO--UR/0216/70/000/002/0302/0305

CIRC ACCESSION NO--AP0117711

UNCLASSIFIED

2/2 010

UNCLASSIFIED

PROCESSING DATE--090CT70

CIRC ACCESSION NO--AP0117711

ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. IT WAS FOUND THAT ILLUMINATION OF YOUNG KIDNEY BEAN PLANTS ENHANCES ACCUMULATION OF AMINOACIDS AND KETOACIDS IN THE LEAVES. THE KETO AND AMINOACIDS CONTENTS IN THE LEAVES IS HIGHER THAN IN THE STEMS AND ROOTS. ILLUMINATION OF THE PLANTS CONSIDERABLY INCREASES INCORPORATION OF C PRIME14 FROM C PRIME14 O SUB2 INTO THE KETOACIDS. NITROGEN STARVATION OF PLANTS ENHANCES THE ACCUMULATION OF KETOACIDS IN THE LEAVES. UTILIZATION OF KETOACIDS FOR AMINOACID SYNTHESIS IN THE LEAVES IS MORE ACTIVE ON AMMONIA NITROGEN THAN ON NITRATE NITROGEN. THUS ON AMMONIA NITROGEN THE KETOACID CONTENTS IN THE LEAVES IS LOWER THAN ON OTHER SOURCES OF NITROGEN.

FACILITY: A. N. BACH INSTITUTE OF BIOCHEMISTRY ACADEMY OF SCIENCES USSR.  
FACILITY: TECHNOLOGICAL INSTITUTE OF FOOD INDUSTRY, MOSCOW.

UNCLASSIFIED

1/2 014 UNCLASSIFIED PROCESSING DATE--20NOV70  
TITLE--EFFECT OF GIBBERELIC AND INDOLEACETIC ACIDS ON PLANT MALATE  
DEHYDROGENASE ACTIVITY -J-  
AUTHOR--(03)-KREIGVICH, V.L., SEVERNAYA, T.A., BUTENKO, R.G.

COUNTRY OF INFO--USSR

SOURCE--DOKL. AKAD. NAUK SSSR 1970, 191(2), 474-5

DATE PUBLISHED-----70

SUBJECT AREAS--BIOLOGICAL AND MEDICAL SCIENCES

TOPIC TAGS--GIBBERELIC ACID, INDOLE DERIVATIVE, PLANT PHYSIOLOGY,  
DEHYDROGENASE, ENZYME ACTIVITY, INHIBITION

CONTROL MARKING--NO RESTRICTIONS

DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FRAME--3003/1475

STEP NO--UR/0020/T0/193/002/0476/0475

CIRC ACCESSION NO--ATD17009

2/2 C14

UNCLASSIFIED

PROCESSING DATE--20NOV70

CIRC ACCESSION NO--AT0130404

ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. INDOLE,3,ACETIC ACID AT 10 PRIME  
NEGATIVES-10 PRIME NEGATIVE6 M INHIBITED MALIC DEHYDROGENASE FROM CORN  
ROOTS AND FROM BOTH NATURAL AND CULTURED TOBACCO WITH PARENCHYMA IN  
VITRO. GIBBERELIC ACID HAD NO SUCH EFFECT. FACILITY: INST.  
BIOKHIM. IM. BAKHA. MOSCOW, USSR.

UNCLASSIFIED

USSR


UDC 621.515.592

UFIMTSEV, V.B., SHUMILIN, V.P., KOSTIN, G.V., KRESTOVNIKOV, A.N.

"Concerning The Character Of The Change Of Composition Of The Vapor Phase Above Type A<sup>III</sup>B<sup>V</sup> Compounds"

Sb.nauch.tr.po probl. mikroelektron. Mosk. in-ta elektron. tekhn. (Collection Of Scientific Works On Problems Of Microelectronics. Moscow Institute Of Electronics Technology), 1972, Issue 8, pp 131-136 (from RZh:Elektronika i yeye primeneniye, No 9, Sept 1972, Abstract No 9B100)

Translation: The paper considers the inverse character of the vapor phase during replacement of the condensate of the element B<sup>V</sup> by the condensate of the compounds A<sup>III</sup>B<sup>V</sup>. From thermodynamic data the temperature dependences are calculated of the vapor pressure of 2- and 4-atom molecules of component B<sup>V</sup> above the six compounds: InSb, GaSb, InAs, GaAs, InP, and GaP. Calculations are made for the reactions



1/2



1/2 010 UNCLASSIFIED PROCESSING DATE--160CT70  
TITLE--SATURATED VAPOR PRESSURE OF MERCURY TELLURIDE -U-  
AUTHOR--(04)-LEVITSKAYA, T.O., VANYUKOV, A.V., KRESTOVNIKOV, A.N., BYSTROV,  
V.P.  
COUNTRY OF INFO--USSR  
SOURCE--IZV. AKAD. NAUK SSSR, NEORG. MATER. 1970, 6(3), 559-60  
DATE PUBLISHED-----70  
SUBJECT AREAS--CHEMISTRY  
TOPIC TAGS--MERCURY COMPOUND, TELLURIDE, VAPOR PRESSURE, THERMAL ANALYSIS  
CONTROL MARKING--NO RESTRICTIONS  
DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FRAME--1996/0834 STEP NO--UR/0363/70/006/003/0559/0560  
CIRC ACCESSION NO--AP0119010  
UNCLASSIFIED

2/2 010

UNCLASSIFIED

PROCESSING DATE--16OCT70

CIRC ACCESSION NO--AP0118010

ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. THE SATD. VAPOR PRESSURE FOR HGTE WAS DETD. OVER COMPNS. CORRESPONDING TO THE HG AND TE BOUNDARIES OF THE HOMOGENEITY REGION, USING THE STATIC COMPENSATION METHOD. THE PRESSURE OVER HGTE ENRICHED WITH HG IS CONSIDERABLY HIGHER THAN THE PRESSURE OVER HGTE ENRICHED WITH TE. THE VAPOR PRESSURE ABOVE HGTE SATD. WITH TE REMAINS CONST. AT 460-560DEGREES INDICATING A CONST. COMPN. OF THE SOLID PHASE; ABOVE THIS POINT, IT BEGINS TO INCREASE, ATTESTING TO ENRICHMENT OF THE SOLID PHASE BY TE AT HIGHER TEMPS. THE HEAT OF FORMATION FOR HGTE IS 16.3 KCAL-MOLE AND THE HEAT OF FUSION, 8.5 KCAL-MOLE. THE M.P. OF HG SUBI-X TE WAS 668DEGREESC, WHICH IN GOOD AGREEMENT WITH THE DTA CURVES; THE CORRESPONDING PRESSURE IS 12.5 ATM. FACILITY: MOSK. INST. STALI SPLAVOV, MOSCOW. USSR.

UNCLASSIFIED

Inorganic Compounds

UDC 541.11

K

USSR

UFIMTSEV, V. B., SHUMILIN, V. P., KRESTOVNIKOV, A. N., and VIGDOROVICH, V. N., Moscow Institute of Steels and Alloys

"Dissociation Pressure of Indium and Gallium Phosphides"

Moscow, Zhurnal Fizicheskoy Khimii, Vol 44, No 4, Apr 70, pp 1120-1122

Abstract: An experimental study of the dissociation pressure of indium and gallium phosphides in the solid state using Knudsen's effusion method is described.  $\lg p = -A/T + B$ -type equations were derived and heats of sublimation determined. Tables in the original article show the rated characteristics of the vapor phase for indium phosphide and data on the temperature dependence of the dissociation elasticity of indium and gallium phosphides. Evaluation of the experimental data by the method of least squares yielded equations describing the temperature dependence of phosphorus vapor pressure. The heats of sublimation derived from these equations are:

$\Delta H_{945} = 49.69 \pm 1.675$  (for InP) and  $\Delta H_{1126} = 65.750 \pm 0.819$  kcal/mole (for GaP).

1/1

1/2 034 UNCLASSIFIED PROCESSING DATE--09OCT70  
TITLE--NEUTRAL DEFECTS IN MERCURY TELLURIDE -U-  
AUTHOR--(03)-LEVITSKAYA, T.O., VANYUKOV, A.V., KRESTOVNIKOV, A.N.  
COUNTRY OF INFO--USSR R  
SOURCE--IZV. AKAD. NAUK SSSR, NEORG. MATER. 1970, 6(3), 556-8  
DATE PUBLISHED-----70  
  
SUBJECT AREAS--PHYSICS, MATERIALS  
TOPIC TAGS--SINGLE CRYSTAL, CRYSTAL LATTICE DEFECT, HALL EFFECT, THERMAL  
EFFECT, IONIZATION, CHEMICAL BONDING, MERCURY COMPOUND, TELLURIDE,  
CRYSTAL GROWTH  
  
CONTROL MARKING--NO RESTRICTIONS  
DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FRAME--1994/1889 STEP NO--UR/0363/70/006/003/0556/0558  
CIRC ACCESSION NO--AP0115708  
UNCLASSIFIED

2/2

034

UNCLASSIFIED

PROCESSING DATE--09OCT70

CIRC ACCESSION NO--AP0115708

ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. A STUDY OF THE KINETICS OF ANNEALING OF HGTE SINGLE CRYSTALS GROWN BY THE BRIDGMAN TECHNIQUE INDICATES THAT THEY CONTAIN NEUTRAL DEFECTS. THIS IS ILLUSTRATED WELL BY COMPARING THE CHANGE IN THE TEMP. DEPENDENCE OF THE HALL COEFF. AND THE CURRENT CARRIER MOBILITIES AS A FUNCTION OF THE ANNEALING TIME AT VARIOUS TEMPS. THE OBSD. RAPID CHANGE IN THE PROPERTIES AT 204 AND 300DEGREES CAN BE CAUSED ONLY BY THE IONIZATION OF NEUTRAL DEFECTS ALREADY EXISTING IN THE CRYSTALS. DEFECTS THAT ARE NEUTRAL AT 204DEGREES BECOME IONIZED AT 358.5DEGREES, WHICH IS REFLECTED IN THE HALL CONST. AND THE CARRIER MOBILITY VALUES. SHORT TERM HIGH TEMP. TREATMENT OF SAMPLES ANNEALED AT 358.5DEGREES DOES NOT CHANGE THE PROPERTIES OF THE HGTE. THUS, NEUTRAL DEFECTS ARE PRESENT IN HGTE AFTER CRYSTN. BY THE BRIDGMAN METHOD, WHICH INHIBIT THE DIFFUSION OF HG INTO THE CRYSTAL WHICH REMAINS NEUTRAL TO SIMILAR TO 360DEGREES BUT IONIZES AT HIGHER TEMPS. INASMUCH AS IONIZATION OF THESE DEFECTS LEADS TO AN INCREASE IN THE ACCEPTOR CONC., THEY APPARENTLY REPRESENT EXCESS TE ATOMS. MORE LIKELY THAN NOT, THIS TYPE OF DEFECT IS ASSOC. WITH THE CHANGE IN THE POSITION OF EXCESS TE IN THE LATTICE, POSSIBLY WITH THE EMERGENCE OF TE-TE BONDS. FACILITY: MOSK. INST. STALI SPLAVOV, MOSCOW, USSR.

UNCLASSIFIED

Semiconductor Technology

USSR

UDC 537.311.33:536.7

VIGDOROVICH, V. N., ~~KRESTOVNIKOV, A. N.~~ and UFINTSEV, V. B., Moscow Institute of Electronic Engineering, Moscow Institute of Steel and Alloys  
Moscow, Ministry of Higher and Secondary Specialized Education USSR  
"Questions of the Topology and Thermodynamics of p-T-X Constitution  
Diagrams of Semiconductor Systems"

Moscow, Izvestiya Akademii Nauk SSSR -- Neorganicheskiye Materialy, Vol 6, No 4, Apr 70, pp 708-713

Abstract: The article considers results of a generalized theoretical analysis of heterogeneous equilibria in semiconductor systems. The heterogeneous equilibria of decomposing semiconductor compounds and phase equilibria at the point of congruent crystallization of semiconductor compounds are considered. It is shown that the theory of regular solutions is applicable to an analysis of the curves for the three-phase equilibrium:  $AB_{sol.} \rightleftharpoons (solution A and B)_{liq.} \rightleftharpoons component B_{vap.}$

The thermodynamics of phase equilibria in systems with continuous series of solid and liquid solutions are considered, as well as changes in vapor-phase composition in the replacement of condensate  $B_{sol.}$  by condensate  $AB_{sol.}$

1/1

1/2 029 UNCLASSIFIED PROCESSING DATE--2009/7/0  
TITLE--TOPOLOGY AND THERMODYNAMICS OF THE P,T,X DIAGRAMS OF SEMICONDUCTING  
SYSTEMS -U-  
AUTHOR-(08)-VIGDOROVICH, V.N., KRESTOVNIKOV, A.M., DEKHTSEV, V.B.

COUNTRY OF INFO--USSR

SOURCE--IZVEST. AKADEMIY NAUK SSSR, NEDRUG. MATERIALY, APR. 1970, 6, (6),  
708-13  
DATE PUBLISHED-----70

SUBJECT AREAS--PHYSICS

TOPIC TAGS--THEORETIC PHYSICS, THERMODYNAMIC ANALYSIS, GALLIUM ARSENIDE  
SEMICONDUCTOR, INDIUM ARSENIDE SEMICONDUCTOR

CONTROL MARKING--NO RESTRICTIONS

DOCUMENT CLASS--UNCLASSIFIED  
PROXY REFERENCE--300470913

STEP 00--0070163770700070057070470713

CIRC ACCESSION NO--AP0101000

2/2 029

UNCLASSIFIED

PROCESSING DATE--20010720

CIRC ACCESSION NO--AP0131499

ABSTRACT/EXTRACT--(U) OP-- ABSTRACT. A GENERAL THEORETICAL ANALYSIS OF THE TEMP., PRESSURE, COMPOSITION DIAGRAMS OF SEMICONDUCTING SYSTEMS OF THE A PRIMEIV A PRIMEIV AND A PRIMEIII B PRIMEV-A PRIMEIII B PRIMEV TYPES SUCH AS GaAs-InAs AND ITS ANALOGUES IS PRESENTED. THE THEORY OF REGULAR SOLUTIONS MAY BE VALIDLY APPLIED TO THE ANALYSIS OF THE THREE PHASE EQUILIBRIUM CURVES OF THESE SYSTEMS. PHASE EQUILIBRIA IN SYSTEMS HAVING A CONTINUOUS SERIES OF SOLID AND LIQUID SOLUTIONS ARE DISCUSSED FROM THE THERMODYNAMIC POINT OF VIEW.

UNCLASSIFIED



1/2 018 UNCLASSIFIED PROCESSING DATE--30OCT70  
TITLE--DISSOCIATION PRESSURE FOR INDIUM PHOSPHIDE AND GALLIUM PHOSPHIDE  
-U-  
AUTHOR--(04)--UFIMTSEV, V.B., SHUMILIN, V.P., KRESTOVNIKOV, A.N.,  
VIGDOROVICH, V.N.  
COUNTRY OF INFO--USSR  
SOURCE--Zh. Fiz. Khim. 1970, 44(4), 1120-2  
DATE PUBLISHED-----70  
SUBJECT AREAS--CHEMISTRY  
TOPIC TAGS--INDIUM COMPOUND, PHOSPHIDE, DISSOCIATION, PRESSURE, GALLIUM  
COMPOUND  
CONTROL MARKING--NO RESTRICTIONS  
DOCUMENT CLASS--UNCLASSIFIED  
PROXY REEL/FRA--1999/1171 STEP NO--UR/0076/70/044/004/1120/1122  
CIRC ACCESSION NO--AP0123148  
UNCLASSIFIED

2/2 018

UNCLASSIFIED

PROCESSING DATE--300CT7C

CIRC ACCESSION NU--AP0123148

ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. THE DISSOCN. PRESSURES OF INP AND GAP WERE STUDIED BY THE KNUDSEN EFFUSION METHOD AT T 879-1011DEGREES K FOR INP AND 1030-1223DEGREESK FOR GAP. THE RESULTS ARE TABULATED. APPLICATION OF THE LEAST SQUARE METHOD TO THE EXPTL. DATA PRODUCED THE FOLLOWING EQUATIONS THAT DESCRIBE THE TEMP. DEPENDENCE OF P VAPOR PRESSURE OVER SOLID INP AND GAP:  $\log P$  (MM HG) EQUALS (MINUS 10,859 PLUS OR MINUS 366-T) PLUS (9.9241 PLUS OR MINUS 0.0699) FOR INP;  $\log P$  (MM HG) EQUALS (MINUS 14,150 PLUS OR MINUS 179-T) PLUS (11.0000 PLUS OR MINUS 0.0722) FOR GAP. THE HEAT OF SUBLIMATION DERIVED FROM THESE EQUATIONS IS  $\Delta H_{\text{SUB}}$  945DEGREES EQUALS 49.691 PLUS OR MINUS 1.675 FOR INP AND  $\Delta H_{\text{SUB}}$  1126DEGREES EQUALS 65.750 PLUS OR MINUS 0.819 KCAL-MOLE FOR GAP. FACILITY: MOSK. INST. STALI SPLAVOV, MOSCON, USSR.

UNCLASSIFIED

1/2 026 UNCLASSIFIED PROCESSING DATE--27NOV70  
TITLE--INVERSION OF VAPOR PHASE COMPOSITION OVER A PRIMEIII B PRIMEV TYPE  
SEMICONDUCTOR COMPOUNDS -U-  
AUTHOR--(04)-UFIMTSEV, V.B., KRESTOVNIKOV, A.N., KOSTIN, G.V., SHUMILIN,  
V.P.  
COUNTRY OF INFO--USSR  
SOURCE--ZH. FIZ. KHIM. 1970, 44(5), 1360  
DATE PUBLISHED-----70

SUBJECT AREAS--PHYSICS

TOPIC TAGS--SEMICONDUCTOR MATERIAL, INDIUM, GALLIUM, ANTIMONY, ARSENIC,  
PHOSPHORUS, VAPOR STATE, PHASE COMPOSITION

CONTROL MARKING--NO RESTRICTIONS

DOCUMENT CLASS--UNCLASSIFIED

PROXY REEL/FRA--3007/0890

STEP NO--UR/0076/10/044/005/1360/1360

CIRC ACCESSION NO--AP0136324

UNCLASSIFIED

UNCLASSIFIED

PROCESSING DATE--27NOV70

2/2 026

CIRC ACCESSION NO--AP0136324

ABSTRACT/EXTRACT--(U) GP-0- ABSTRACT. EQUIL. VAPOR PRESSURES OF B SUB2  
PRIMEV AND B SUB4 PRIMEV OVER A PRIMEIII B PRIMEV SOLIDS WERE CALCD.  
FROM KNOWN THERMODYNAMIC PROPERTIES AND CORRELATED BY THE EQUATION  $\log P$   
EQUALS MINUS (A-T) PLUS B FOR A PRIMEIII EQUALS IN OR GA AND B PRIMEV  
EQUALS SB, AS, OR P. THE VALUES OF A ARE GREATER FOR B SUB4 THAN FOR B  
SUB2 OVER A PRIMEIII B PRIMEV WHILE WITH PURE SOLID B, THE REVERSE IS  
TRUE. FACILITY: MOSK. TEKHNOL. INST. STALI SPLAYOV, MOSCOW,  
USSR.

UNCLASSIFIED